



## **Tramadol Induced Ovarian and Uterine Changes in Albino Rats**

**Girish Rathod <sup>a\*</sup>, Somanath Reddy Patil <sup>b</sup>, Md. Liyakat Ahmed <sup>c</sup>  
and K. Vijaykumar <sup>a</sup>**

<sup>a</sup> Department of PG Studies and Research in Zoology, Gulbarga University, Kalaburagi- 585106, Karnataka, India.

<sup>b</sup> Department of PG Studies and Research in Zoology, Government (Autonomous) College, Kalaburagi- 585106, Karnataka, India.

<sup>c</sup> Department of Pharmacology, Luqman College of Pharmacy, Old Jewargi Road Kalaburagi- 585102, Karnataka, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JPRI/2021/v33i51B33535

Editor(s):

(1) Dr. Takashi Ikeno, National Institute of Mental Health, National Center of Neurology and Psychiatry, Japan.

Reviewers:

(1) Rafael Alberto González Cortés, Latin University of Costa Rica, Costa Rica.

(2) Laurentiu Simion, University of Medicine and Pharmacy Carol Davila, Romania.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available in this link:

<https://www.sdiarticle5.com/review-history/77338>

**Original Research Article**

**Received 09 September 2021**

**Accepted 19 November 2021**

**Published 26 November 2021**

### **ABSTRACT**

Tramadol at the dose levels of 1mg and 3mg/100g body weight was administered to normal cycling rats for 20 days through intraperitoneal routes. At autopsy on 21<sup>st</sup> day significant reduction in the ovarian, uterine and body weight was observed. Histological observations showed decrease in the number and size of Graafian follicles, corpora lutea and increase in the atretic follicles in the ovary. The uterus showed absences of endometrial glands, decrease in the height of myometrium, endometrium and its epithelial cells. The total protein and glycogen content of the ovary and uterus is decreased whereas the cholesterol content is increased. The hypothalamo-hypophyseal gonadal axis is prominent regulator of reproductive activities in animals through neuro-endocrine regulation. In this study action of tramadol on ovary and uterine parameters is discussed.

**Keywords:** Tramadol; atretic follicles; ovarian steroids; neuro-endocrine.

## 1. INTRODUCTION

Tramadol is an analgesic with opioid-like effects [1] and [2]. It has pharmacodynamic and pharmacokinetic profile relative to other opioids. The treatment of mild to severe pain at dose of 200 or 300 milligrams per day is the formulations in both immediate and extended-release in many parts of the world [3] and [4].

The toxicity of tramadol includes central nervous system depression, nausea, vomiting, dizziness, anorexia, seizures and hypotension which may occur in therapeutic or toxic doses [5]. Fatal toxicity of tramadol have been reported as a result of overdose. In these cases, death has been result due to cardiopulmonary arrest and liver failure, in addition hypoglycaemia [6] and [7].

In the year 2013 more than 44 million people were prescribed tramadol in United States, making it one of the most used opioids [8] Many countries of the world use moderate to severe medications of opioid-based analgesics. During the last 20 years, there is significant increase and also higher demand of opioids globally [9]. The acceptability and availability of opioids for the treatment of pain differs throughout the world [10] and [11].

As over development or reproductive toxicity, endogenous opioid peptides are said to be located in different tissues on the reproductive system which indicate that they might be involved in the reproductive functions [12,13].

As peptides induces their effect on opioid receptors. So, in men it can cause loss of libido and erectile and ejaculatory dysfunctions and can inhibit the proliferation of uterine cells which mediate mainly by mu opiate receptor [14]. Chronic administration of tramadol can cause reproductive dysfunction and increased average of sterility [15]. The abuse of opiate leads to hypogonadism, primarily by decreasing the release of gonadotropin releasing hormone(GnRH), deficiency of testosterone and infertility. Many researchers have demonstrated that long-term administration of tramadol had dose dependent adverse effects on testicular tissue [16]. Some other studies have showed that rats received subcutaneous injections of tramadol (40 mg/kg body weight) three times per week for 8 weeks showed reduced plasma levels

of luteinizing hormone, follicle stimulating hormone [17,18]. In addition, tramadol caused a concentration-dependent inhibition of potassium chloride-induced myometrium contractility.

## 2. MATERIALS AND METHODS

Normal cycling healthy female albino rats of wistar strain were maintained at room temperature of  $28 \pm 2^\circ\text{C}$  with lighting schedule of 12 hours light and 12 hours darkness. They were grouped in individual cages, each containing six animals. They were fed with a standard pellet diet (VRK Nutrition, Pune) and water *ad libitum*.

The animals were divided into the following groups:

Group 1- Received 0.2ml saline interaperitonally for 20 days and served as control group.

Group 2- These rats received tramadol 1mg/100g body weight interaperitonally for 20 days.

Group 3- These rats received tramadol 3mg/100g body weight interaperitonally for 20 days.

The treatment was started from estrous phase of the cycle only as the ovarian and uterine activities change markedly from one phase to another phase. The treatment was given once a day between 10:00 AM to 11:00 AM for 20 days. All the experimental rats were sacrificed by decapitation on 21<sup>st</sup> day 24 hours after the final dose.

The body weight was recorded. Ovary and Uterus were dissected out, freed from adherent tissue and weighed on Anamed electronic balance. The number of Graafian follicle, atretic follicle and corpora lutea was made from randomly chosen 20 sections from each group. Micrometric measurements such as diameter of uterus, thickness of myometrium, endometrium and epithelial cell height were also made from randomly selected 20 sections which appeared round incross section from each group. Micrometric measurements were made by using stage and ocular micrometer.

Protein content of ovary and uterus was estimated by Lowry's method [19]. Cholesterol content was done by Libermann and Burchard's reaction as described by Peter and Vanslyke [20]. The glycogen content of ovary and uterus

was estimated by Carrol *et al.*, Statistical analysis was carried out by using student “t” test.

### 3. RESULTS

Body weight – There is no significant change in the body weight of the rats due to treatment of tramadol for 20 days, intraperitoneally compared to their respective control group.

In comparison with the control group of rats highly significant ( $P<0.01$ ) increase in the cholesterol content and significant decrease ( $P<0.001$ ) in protein and glycogen content of the ovary was observed in the treated group of rats.

Due to the administration of tramadol, the number of healthy follicles is decreased in both 1mg and 3 mg showing significant decrease in class I to class VI follicles.

In comparison with the control group of rats highly significant ( $P<0.01$ ) increase in the cholesterol content and significant decrease ( $P<0.001$ ) in protein and glycogen content of the uterus was observed in the treated group of rats.

There was significant reduction in the diameter of uterus, thickness of endometrium and

myometrium and epithelial cell height ( $P<0.01$ ) in tramadol treated groups compared to their respective control group. A reduction in the secretion of endometrial gland was observed.

### 4. DISCUSSION

It is also known that hypothalamus regulates the rhythmic release of pituitary gonadotrophins, i.e., FSH, LH and prolactin through neural stimulus to GnRH [21]. The orderly event of follicular growth and ovulation depends upon the pituitary FSH, LH and prolactin. Investigations on tramadol indicate that tramadol being a central nervous system influencing drug inhibits the release of gonadotrophins from pituitary [22]. The studies also indicate that tramadol blocks ovulation by inhibiting the LH surge from pituitary in rats [23,24]. In the present study, as the drug was administered between 10.00 and 11.00 AM every day, it covers the “critical period” of LH surge, thus postponing the ovulation for one day by interfering with 24 hours periodicity for gonadotrophin release. Decreased level of protein content in tramadol treated rats indicates hampered growth and source of energy. Increased levels of cholesterol and glycogen content indicated hampered steroidogenesis.

**Table 1. Effect of Tramadol on gravimetric and biochemical changes of ovary in albino rat**

	Weight (mg/100g body wt.) Ovary	Cholesterol ( $\mu$ g/mg) Ovary	Protein ( $\mu$ g/mg) Ovary	Glycogen ( $\mu$ g/mg) Ovary
Saline	46.51 $\pm$ 0.94	3.23 $\pm$ 0.03	6.01 $\pm$ 0.05	2.43 $\pm$ 0.03
1mgTramadol	43.40 $\pm$ 0.69**	4.41 $\pm$ 0.05**	5.16 $\pm$ 0.18**	1.71 $\pm$ 0.05**
3mgTramadol	40.66 $\pm$ 1.17**	5.63 $\pm$ 0.07**	4.31 $\pm$ 0.06**	1.54 $\pm$ 0.06**

*M $\pm$ S = Mean  $\pm$  Standard Error*

*\*P<0.01; \*\*P<0.001, compared to respective control.*

**Table 2. Effect of Tramadol on Ovarian components in albino rats**

Treatment	Class I SPAF	Class II LPAF	Class III SAF	Class IV MSAF	Class V LSAF	Class VI GF
Saline	58.61 $\pm$ 3.1	49.17 $\pm$ 2.17	11.02 $\pm$ 0.76	5.18 $\pm$ 0.81	3.38 $\pm$ 0.03	2.46 $\pm$ 0.37
1mgTramadol	34.26 $\pm$ 1.33**	31.89 $\pm$ 0.40**	7.82 $\pm$ 0.31**	4.61 $\pm$ 0.08**	2.23 $\pm$ 0.33**	1.01 $\pm$ 0.02**
3mgTramadol	29.38 $\pm$ 0.51**	28.51 $\pm$ 1.12**	7.40 $\pm$ 0.03**	3.44 $\pm$ 0.04**	1.17 $\pm$ 0.19**	0.68 $\pm$ 0.01**

*M $\pm$ S = Mean  $\pm$  Standard Error*

*\*P<0.01; \*\*P<0.001, compared to respective control*

**Table 3. Effect of Tramadol on gravimetric and biochemical changes of uterus in albino rats**

	<b>Weight (mg/100g body wt.) Uterus</b>	<b>Cholesterol (µg/mg) Uterus</b>	<b>Protein (µg/mg) Uterus</b>	<b>Glycogen (µg/mg) Uterus</b>
Saline	228.21±4.21	4.18±0.02	8.28±0.09	1.64±0.05
1mgTramadol	201.81±3.61*	5.51±0.14**	5.16±0.02**	1.49±0.07*
3mgTramadol	169.67±2.28**	6.75±0.05**	4.37±0.03**	1.31±0.04**

*M±S = Mean ± Standard Error*

\**P*<0.01; \*\**P*<0.001, compared to respective control

**Table 4. Effect of Tramadol on histometric changes of uterus in albino rat**

	<b>Diameter of uterus (µm)</b>	<b>Thickness of myometrium (µm)</b>	<b>Thickness of endometrium(µm)</b>	<b>Height of epithelium (µm)</b>
Saline	2139.49±7.38	241.01±2.29	433.00±2.08	39.89±1.02
1mgTramadol	1914±8.23**	183.01±2.04**	349.18±3.49**	29.28±2.12**
3mgTramadol	1737.36±9.27**	170.17±3.21**	317.59±5.89**	19.24±0.8**

*M±S = Mean ± Standard Error*

\**P*<0.01; \*\**P*<0.001, compared to respective control

FSH stimulates the differentiation of granulosa cells and promotes the follicular development [25,26]. In the present investigation, there is reduction in the number of Graffian follicle in the ovary of tramadol treated rat, thus indicating the inhibition of follicular growth which is gonadotrophin dependent. There is decrease in the number of corpora lutea in tramadol treated rat indicating reduction in the rate of ovulation leading to follicular atresia.

The growth of uterine cavity depends upon the ovarian estrogen secretion. Primarily estrogen acts upon the surface epithelium and the glands within endometrium [27]. Progesterone acts on estrogen primed uterus and thus prepares the uterine epithelium from proliferative to secretory state.

## 5. CONCLUSION

In the present study as there is no significant change in the body weight of albino rats was observed indicates that tramadol has no effect on body weight. Decreased ovarian weight along with protein and glycogen content shows that hampered physiological activity and source of energy.

Decreased in the ovarian components indicates that ovarian activity has been hampered and that may be attributed to non-availability of hormones. Reduction in the diameter of uterus, thickness of endometrium, myometrium and epithelial cell height may be attributed to the non-availability of

steroids necessary for uterine growth due to decreased levels of gonadotrophins in tramadol treated rats.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

Approval at the Institutional Animal Ethics Committee (IAEC) of Luqman College of Pharmacy, Gulbarga was taken for conducting experimental activities.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Volk R, Melhuish T, Chong C, Ryan T, White LD. Adjuncts to local anaesthetics in tonsillectomy: a systematic review and meta-analysis. *J Anesthesia*. 2017;31(4): 608-16
2. Schnabel A, Reichl SU, Meyer-Friebem C, Zahn PK, Pogatzki-Zahn E. Tramadol for postoperative pain treatment in children. *Cochrane Database Syst Rev*. 2015;18(3)
3. Vijayan R, Afshan G, Bashir K, Cardosa M, Chadha M, Chaudakshetrin P, et al.,

- Tramadol: a valuable treatment for pain in Southeast Asian countries. *J Pain Res.* 2018;(11):2567-75
4. Santos Garcia JB, Lech O, Campos Kraychete D, Rico MA, Hernandez-castro JJ, Colimon F, *et al.* The role of tramadol in pain management in Latin America: A report by the change pain Latin America advisory panel. *Curr Med Res Opin.* 2017; (9):1615-21
  5. Beakley BD, Kaye AM, Kaye AD. Tramadol, pharmacology, side effects and serotonin syndrome: A review, *PainPhysician.* 2015;18(4):395-400
  6. Daubin C, Quentin C, Goulle JP, Guillotin D, Lepage O. *Clin Toxicol (phila).* 2007 (45)961
  7. Mugunthan, N, Davoren P. *Endocrine Practical.* 2012;18:151.
  8. Patterson E. *Tramadol History and Statistics;* 2017
  9. Berterame S, Erthal J, Thomas J, Fellner S, Vosse B, Clare P, *et al.*, Use of and barriers to access to opioid analgesics: A worldwide, regional and national study. *Lancet.* 2016; 387(10028):1644-56.
  10. Kunumpurath S, Julien N, Kodumudi G, Kunumpurath A, Kodumudi V, Vadivelu N. Global supply and demand of opioids for pain management. *Curr Pain Headache Rep.* 2018;22(5):34.
  11. Pastrana T, Wenk R, Radbruch L, Ahmed E, De Lima L. Pain treatment continues to be inaccessible for many patients around the globe: second phase of opioid price watch, a cross-sectional study to monitor the prices of opioids. *J Palliative Med.* 2017;20(4):378-87
  12. Subiran N, Casis L, Irazusta JR, Regulation of Male Fertilty By the Opioid System. *Molecular Med.* 2011;17(7-8); 846-853
  13. Vuong C, Van Uvm S, O'dell L, Lutfy K and Friedman T. The effect of opioids and opioid analogs on animal and human endocrine systems. *Endocrine Rev.* 2010;31(1):98-132
  14. Dziekonski M, Zmijewska A, Franczak A, Kotwica G, Kosciukiewicska W, Okrasa S. The expression of mRNA's for opioid receptors in endometrium of cyclic and early pregnant pigs; in-vitro effects of IL-1 $\beta$ , IL-6 and TNF  $\alpha$ (alpha). *Journal of Animal and fed Science.* 2015;24(4):332-340
  15. El-Ghawet HA. Effects of tramadol on the reproductive function of wister albino rats. *Eur J Exp Biol.* 2015;5:56-64
  16. Azari O, Emadi L, Kheirandish R, Shafiei Bafti H, Esmali Nejad MR, Farghi F. The effects of long term administration of tramadol on epididymal sperm quality and testicular tissue in mice. *International Journal of Veter Sur.* 2014;9(1):23-30
  17. Ahmed MA, Kurkar A. Effects of opioid (tramadol) treatment on testicular functions in adult male rats. The role of nitric oxide and oxidative stress. *Clinical and Experimental Pharma and Phys.* 2014;41(4):317-323
  18. Vazzana M, Andreani T, Fangueiro J, Faggio C, Silva, Santini A, Garcia M, A, Souto E. Tramadol hydrochloride: Phaqrmacokinetics, Pharmacodynamics; adverse side effects, co-administration of drugs and new drugs delivery systems. *Bio-Med & Pharma.* 2015;70:234-238
  19. Lowry OH, Rosenbrough NJ, Farr NL, Randoll RJ, Protein measurement with folin. Phenol reagent. *J Biol Chem.* 1951;193:265-275.
  20. Peter's JP, Vanslyke DD, *Quantitiative Clinical Chemistry Vol. I, Williams and Wilkins.*(Eds), Baltimore.1946.
  21. Carmel PW, Arakis, Ferin M. Pituitary stalk portal blood collection in Rhesus Monkeys: Evidence or pulsatile release of gonadotrophin releasing hormone(GnRH). *Endocrinology.* 1976;99:243-248.
  22. Salah S, Wagih M, Zaki A, *et al.*, Long term effects of tramadol on the reproductive function of male albino rats: an experimental biochemical and histopathological study. *Middle East Fertil Soc J.* 2020;24:3
  23. Lawton I, Sawyer CH. Timing of gonadotrophin and steroid secretion at diestrus in the rat. *Endocrinology.* 1968;83:831-836.
  24. Sindgi SB. Effect of barbiturates on ovarian growth and pregnancy in albino rats. Ph.D. Thesis, Karnataka University, Dharwad, India. 1975.
  25. Channing CP. Influences of the *vivo* and *vitro* hormonal environment. *Recent Prog Horm Res.* 1970;26:589-593.
  26. Goldenberg RL, Vaitukaitis JL, Ross STG. Estrogen and follicle stimulating hormone interactions on follicle growth in rats. *Endocrinology.* 1972;90:1492-1496.

27. Richards JS, Ireland JJ, Rao MC, Benath GA, Midgley Jr. AR, Reichert LE. Ovarian follicular development in rat hormone receptors regulation by Oestrodiol, follicle stimulating hormone and luteinizing hormone. *Endocrinology*. 1976;87:330-334.

---

© 2021 Rathod et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/77338>