



# Well Differentiated Villoglandular Papillary Adenocarcinoma of the Cervix: A Rare Case Report

Anita Sajjanar <sup>a≡\*</sup>, Pratibha Dawande <sup>a<sup>o</sup></sup>, Swetha Bahadure <sup>a≡</sup>  
and Sunita J. Vagha <sup>b<sup>o</sup></sup>

<sup>a</sup> Department of Pathology, Datta Meghe Medical College, Shalinitai Meghe Hospital and Research Centre, Nagpur-441110, India.

<sup>b</sup> Department of Pathology, Jawaharlal Nehru Medical College, Datta Meghe Institute of Medical Sciences, Sawangi (Meghe), India.

## Authors' contributions

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## Article Information

DOI: 10.9734/JPRI/2021/v33i60B35085

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/80108>

**Case Report**

**Received 20 October 2021**  
**Accepted 26 December 2021**  
**Published 28 December 2021**

## ABSTRACT

The Villoglandular papillary adenocarcinoma (VGA) is a variant of adenocarcinoma of the cervix. It occurs in young women having a favorable prognosis. Pathologically, VGA poses a diagnostic challenge. Here we present a rare case of a middle-aged female patient complaining of bleeding per vagina since 1 year. Histopathological examination reported it as a well-differentiated villoglandular adenocarcinoma of the cervix. This tumor as a separate entity should be recognised, distinguished from other variants of adenocarcinomas. Because of its good behaviour and favourable prognosis usually it allows conservative treatment.

**Keywords:** Well-differentiated; villoglandular; papillary; adenocarcinoma; cervix.

## 1. INTRODUCTION

Adenocarcinoma is the second most frequent carcinoma of the cervix [1]. The villoglandular

papillary adenocarcinoma (VGA) is a variant of adenocarcinoma of cervix. First it was described by Young and Scully in 1989 [2,3]. In the review of literature, 56 cases have been reported. This

<sup>≡</sup> Associate Professor;

<sup>o</sup> Professor and HOD;

\*Corresponding author: E-mail: anitavijay28@gmail.com;

tumor was included with other varieties of adenocarcinomas of the cervix and not as a separate entity in the past. Villoglandularpapillary adenocarcinoma of the cervix has favourable prognosis. When the tumor is superficial without vascular or lymphatic invasion, it can be treated by a radical hysterectomy.

## 2. CASE REPORT

A 46-year-old para 2 presented with a history of bleeding per vagina for three months. On per speculum examination, exophytic growth protruding from the cervix was noted for which a punch biopsy was done. The histopathological examination showed well villoglandular differentiated papillary adenocarcinoma. Later she underwent total radical hysterectomy and infracolic omentum. External surface of cervix shows grey, white to brown exophytic polypoid growth with papillary excrescences measuring 6.5x5x4cm (Fig. 1). Cut section show growth


involving cervix (Fig. 2). Cut section of uterus show endometrial thickness is 0.2cm and myometrial thickness is 2cm. Cut section of both ovaries and fallopian tubes were normal. Microscopy examination from growth of cervix show tumor cells arranged in the form of villoglandular and papillary structures with fibrovascular core infiltrated by inflammatory cells (Fig. 3, Fig. 4). These tumor cells are lined by columnar cells having stratified hyperchromatic nuclei with moderate cytoplasm and numerous mitotic figures. Adjacent area shows haemorrhage and necrosis. Endometrium, isthmus, vagina, and omentum is free from invasion. The right and left parametrium shows tumor infiltration. Both, ovary and fallopian tubes were free from tumor deposits [4,5] isolated left and right external iliac lymph nodes show reactive changes and free from tumor deposits. Finally reported as well-differentiated villoglandular papillary adenocarcinoma cervix (FIGO stage IIb) (TNM stage- T1IbN0M0).




**Fig. 1. External surface of cervix shows grey, white to brown exophytic polypoid growth with papillary excrescences along with omentum and pelvic lymph nodes**



**Fig. 2. Cut section show growth involving cervix**



**Fig. 3. Microscopic examination from the growth of the cervix shows tumor cells arranged in villoglandular and papillary structures with fibrovascular core infiltrated by inflammatory cells**



**Fig. 4. Tumor cells are lined by columnar cells having stratified hyperchromatic nuclei with moderate cytoplasm and numerous mitotic figures**

### 3. DISCUSSION

The cervical adenocarcinoma incidence is increasing. It accounts for one-fourth of all cervical carcinomas. Among these VGA accounts for approximately 4% of all cervical adenocarcinomas [6]. Adenocarcinoma usually affects women in their second to fourth decade. Aetiology of VGA is ambiguous. Many studies have found that underlying HPV infection might have been the causative agent of VGA. The first reported positive HPV-18 in VGA was published in year 2000 [7]. Usually, patients present with vaginal bleeding or pelvic pain. Grossly VGA

presents exophytic polypoid friable growth protruding from the cervix. Histologically it is characterized by tall and thin papillae lined by endocervical, endometrial or intestinal-type epithelium with mild atypia [8]. The characteristic histological features of various epithelia have been reported, and 1 or more layers of cells line papillae with endocervical, endometrioid, or intestinal differentiation. VGA are diagnosed and managed successfully with conservative measures. Radical hysterectomy is an adequate treatment for this variant of adenocarcinoma. The prognostic factors are tumour size, tumour grade and clinical stage. Wang et. al. concluded that

stage is the most important prognostic factor with 5-year overall survival rates varying on stage: FIGO stage I - 79%, II - 37%, III / IV - less than 9% [9]. Studies reflecting on different types of cervical cancer [10-12], ovarian tumours [13-16] were reviewed.

#### 4. CONCLUSION

Clinician and pathologist needs to be aware this rare malignancy from uterine cervix. This tumor as a separate entity should be recognized, distinguished from other variants of adenocarcinomas. Because of its good behaviour and favourable prognosis usually it allows conservative treatment. This case is a confirmed well-differentiated VGA of the cervix characterized by an exophytic polypoid growth pattern with villoglandular and papillary structures with a fibrovascular core. These tumor cells are lined by columnar cells having stratified hyperchromatic nuclei with moderate cytoplasm and numerous mitotic figures. Recurrence rate of VGA's compared to other variants of adenocarcinoma of cervix is less.

#### CONSENT

We have added the Consent Disclaimer in the revised paper. The revised paper is attached herewith this mail for your kind perusal. Kindly check the revised paper.

#### ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Keiichi Fujiwara Bradley Monk, Mojgan Devouassoux-Shisheboran. Adenocarcinoma of the uterine cervix: why is it different? *Curr Oncol Rep.* 2014;16(12): 416.
2. Utsugi K, Shimizu Y, Akiyama F, Umezawa S, Hasumi K. Clinicopathologic features of villoglandular papillary adenocarcinoma of the uterine cervix. *Gynecol Oncol.* 2004; 92:6470.
3. Young RH, Scully RE. Villoglandular papillary adenocarcinoma of the uterine cervix. A clinicopathologic analysis of 13 cases. *Cancer.* 1989;63: 17713.
4. Datta CK, et al. Well-differentiated papillary villoglandular adenocarcinoma of the uterine cervix *Med J.* 1997;93(4):186-8.
5. Reed W, Abeler VM, Tropé CG Tidsskr Nor Laegeforen. Villous glandular adenocarcinoma of the uterine cervix. A subtype with favourable prognosis? 1993;113(20):2569-71.
6. Wells M, Ostor AG, Crum CP, Franceschi S, Tommasino M, Nesland TM, et al. Tumors of the cervix. In: World Health Organization, editor. *Tumors of the breast and female genital organs.* Lyon: IARC Press. 2003;262276.
7. Yamazawa K, Matsui H, Seki K, et al. Human papillomavirus-positive well-differentiated villoglandular adenocarcinoma of the uterine cervix: a case report and review of the literature. *Gynecol Oncol.* 2000;77(3):473-477.
8. Kurman RJ, Ronnett BM, Sherman ME, Wilkinson EJ. *Atlas of Tumor Pathology: Tumours of the Cervix, Vagina and Vulva.* Washington, DC: American Registry of Pathology. 2010;179-185. (4th Series).
9. Sophia S. Wang, Mark E. Sherman, Allan Hildesheim, James V. Lacey Jr, Susan Devesa. Cervical adenocarcinoma and squamous cell carcinoma incidence trends among white women and black women in the United States for 1976-2000. *Cancer.* 2004;100(5):1035-44.
10. Prasad R, Khan M, Anjum F, Bhojar S, Acharya N. Tuberculosis of uterine cervix presenting with recurrent leucorrhoea in guise of carcinoma of cervix. *Journal of Clinical & Diagnostic Research.* 2019; 13(11). Available: <https://doi.org/10.7860/JCDR/2019/42669.13296>.
11. Shelke UV, Acharya S, Shrivastava DS. Co-Existence of Chronic Lymphocytic Leukaemia and Malignancy of Uterine Cervix. *Journal of Evolution of Medical and Dental Sciences.* 2020;9(18):1522-5. Available: <https://doi.org/10.14260/jemds/2020/332>.
12. Shukla S, Rajput D, Acharya S, Acharya N, Grover S, Samal N, Vagha S. A study on the nucleolar organizer regions-as a tumour proliferative marker in cervical smears. *Journal of clinical and diagnostic research: JCDR.* 2013;7(2):278.

- Available:<https://doi.org/10.7860/JCDR/2013/4808.27406>
13. Choudhary SV, Banode PJ, Bhake A, Dwivedi S, Gupta S, Koley S, Singh A. Hirsutism with virilization in a postmenopausal woman due to a rare ovarian steroid cell tumor. Available:<https://doi.org/10.4103/0378-6323.60546>
  14. Sinha A, Hariharan C, Chaudhary RK. Preoperative evaluation of ovarian tumors in the rural population--clinical features, grey scale ultrasound, colour Doppler, serum CA-125 and Alcazar score in predicting malignancy. Journal of Evolution of Medical and Dental Sciences. 2013; 2(16):2598-610. Available:<https://Doi.Org/10.14260/Jemds/584>
  15. Mounika P, Shrivastava DS, Diwani D. Umbilical Metastasis Secondary to Ovarian Carcinoma--A Rare Case of Sister Mary Joseph Nodule. Journal of Evolution of Medical and Dental Sciences. 2020;9(44): 3343-5. Available:<https://doi.org/10.14260/jemds/2020/734>
  16. Toshniwal SM, Inamdar SA, Agarwal HD, Sharma SR. Malignant brenner tumor-a rare case of ovarian carcinoma. Medical Science. 2020;24(105):3550-4.
  17. Binu IS, Diwagar KN, Sekhar G. A clinicopathological study of malignant tumors of the uterine corpus in a tertiary care center. Journal of Pharmaceutical Research International. 2021;33(49B):89-96. DOI: 10.9734/jpri/2021/v33i49B33343

© 2021 Sajjanar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/80108>