

Nurses Knowledge and Attitudes on Pain Management in Children at the Tamale West Hospital

Nicholas L. Yombe¹, Vida Nyagre Yakong²
and Keren-Happuch Twumasiwaa Boateng^{3*}

¹Department Paediatric Nursing and Child Health, School of Nursing and Midwifery, University for Development Studies, Tamale, Ghana.

²School of Nursing and Midwifery, University for Development Studies, Tamale, Ghana.

³Department Midwifery and Women's Health, School of Nursing and Midwifery, University for Development Studies, Tamale, Ghana.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Asmaa Fathi Moustafa Hamouda, Jazan University, Saudi Arabia.

Reviewers:

(1) Sharareh Zeighami Mohammadi, Islamic Azad University, Iran.

(2) Ashraf Direkvand Moghadam, Ilam University of Medical sciences, Iran.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:
<https://www.sdiarticle5.com/review-history/78229>

Original Research Article

Received 06 October 2021

Accepted 13 December 2021

Published 13 December 2021

ABSTRACT

Background: Pain is one of the worse symptoms experienced by hospitalized patients and the major reason for seeking medical and surgical intervention. Despite the existence of guidelines on pain assessment and management, children continue to experience unrelieved pain during hospitalization.

Purpose: The study aimed at assessing Nurses Knowledge on Paediatric Pain Assessment and Management at the Tamale West Hospital.

Methods: A descriptive cross-sectional survey was conducted among 120 registered nurses at the Tamale West Hospital in Ghana. The consecutive sampling technique was employed to recruit participants. A structured questionnaire was used for data collection. Data was analyzed and presented using descriptive and inferential statistics. Data was collected using the Pediatric Nurses' Knowledge and Attitudes Survey Regarding Pain.

Findings: The study revealed that majority of nurses at the Tamale West Hospital in Ghana have insufficient knowledge toward pain in children. There is the urgent need to intensify education in this area so as to adequately prepare these nurses to cater for the pain needs of vulnerable children and their families.

Recommendations: Future studies should focus on how student's theoretical knowledge and attitudes are linked to pediatric pain assessment and management practices.

Keywords: Nurses knowledge; pain assessment and management; paediatric ward; paediatric nurses; tamale west hospital.

1. INTRODUCTION

Pain is one of the worse symptoms experienced by hospitalized patients and the major reason for seeking medical and surgical intervention [1]. Despite the existence of guidelines on pain assessment and management, children continue to experience unrelieved pain during hospitalization [2,3]. Inadequate pain management in children is a global health concern due to difficulties encountered in assessing and managing pain in this vulnerable group [4]. Infants and young children may not be able to describe the nature, location and severity depending on their age and cognitive development [2,5]. Hence, nurses who form the majority of healthcare providers may be required to use appropriate strategies to assess and manage children's pain based on their level of cognitive development and pain typology [6].

Efficient paediatric pain assessment and management, relies on sufficient knowledge of health care providers particularly nurses, who spend the most time with hospitalized children and their families. The consequences of unrelieved and mismanaged paediatric pain are not limited to sensations of discomfort and unpleasantness but incorporates further insidious body system malfunctioning [7,8]. These include hyperglycemia, increased inflammatory responses, gastrointestinal obstruction, immunosuppression, chronic pain among others [9,10]. Furthermore, it poses financial burden for affected families and healthcare systems due to delayed recovery time and its associated increased cost of healthcare [11]. Unrelieved pain has been noted to significantly affect children's school attendance, performance, and their ability to play and socialize [12].

Considering the effects of unrelieved pain on children's health, it is not surprising that effective pain management has been recognized as a fundamental human right of every child [13]. Efficient pain management, however, partly relies on sufficient knowledge and appropriate attitudes of health care providers particularly nurses, who spend the most time with hospitalized children and their families. As members of the healthcare

team, nurses play a vital role in the management and control of pain through accurate assessment, planning, implementation and evaluation of pain relief measures [14]. In order for nurses to perform their pain assessment and management roles efficiently and effectively, they need to be adequately knowledgeable. Nurses ought to have a solid knowledge base of pediatric pain and its management as this impact on their practice [14]. Data regarding nurses' knowledge on paediatric pain assessment and management is sparse particularly in the sub-Saharan Africa sub-region including Ghana. The study seeks to assess nurses' knowledge of paediatric pain assessment and management at the Tamale West Hospital.

2. MATERIALS AND METHODS

2.1 Study Design

Research design is a map or plan of how the researcher intends to carry out the research (De Vos et al., 2011). A research design focuses on the end product, formulates a research problem as a point of departure, and focuses on the logic pertaining to the research (De Vos et al, 2011). Research designs are made of quantitative, qualitative or mixed methods approach which gives specific directions for research procedures (Creswell, 2014, p.12). Quantitative research involves the use of computational, statistical, and mathematical tools to derive results (Neuman, 2013). It is conclusive in its purpose as it tries to quantify the problem and understand how prevalent it is by looking for projectable results to a larger population (Neuman, 2013). The present study employed descriptive cross-sectional quantitative approach. This design was appropriate as the study was predictive in nature and examined associations (Shaughnessy, Zechmeister, & Zechmeister, 2012).

2.2 Study Population

All nurses working in the Tamale West Hospital constituted the target population for this study. Specialization in children's nursing is not required of nurses to practice within pediatric care settings in Ghana. Many clinical nurses

encountered in children's care unit are general nurses as the pediatric nursing specialty is still new and evolving in Ghana. Nurses who work in other units of the hospital are also confronted with children who have varied levels of pain caused by clinical conditions, injuries, hospital procedures or some unknown factors. Additionally, nurses in the study hospital are reshuffled annually. This implies that every nurse is likely to work in the paediatric unit if they have never worked there.

2.3 Setting

The study was conducted at the Tamale West Hospital (TWH). The TWH is located in the Tamale metropolis, the administrative capital of the northern region of Ghana. The Tamale Metropolis has a total population of 371, 351 (Ghana Statistical Service, 2014). The hospital has a bed capacity of 120 and serves as a referral facility for some districts in the northern region. As at December 2019, the facility had a total number of 350 nurses. The TWH renders outpatient and inpatient services, medical and surgical treatments, obstetric/gynaecological services, public health services, among others. The hospital was established in April, 1998.

2.3.1 Inclusion criteria

Participants were included in the study if they were:

Registered Nurses who were currently working at all the wards and providing care for children at Tamale West Hospital.

Nurses who had been working at the Hospital for the past one year and currently rotated to work at other wards.

Nurses who had completed a three or/and four-year diploma or/and Bachelor (BSc.) of Nursing.

2.3.2 Exclusion criteria

The exclusion criteria include;

Nurse with less than one year working experience and those who refuse to participate were excluded from the study.

Nurse assistants or enrolled nurses were excluded.

Nurses who were undertaking their one-year rotation were also excluded.

2.4 Sampling

The non-probability consecutive sampling technique was employed to recruit respondents into the study. This technique is chosen to enable the researcher enroll every participant until the required sample size is met.

2.5 Sample Size Determination

Sample size is defined as a count of the individual samples or observations in any statistical such as scientific or public opinion survey (Polit & Beck, 2004). The total sample framework for the number of nurses working in the Tamale west hospital is 170. The sample size for the survey was computed according to the formula for sample size determination by Yamane [15].

$$n = \frac{N}{1+Ne^2}$$

Where n is required sample size.

N is the total population size which is 170 nurses.

e is acceptable sampling error (0.05) at 95% Confidence Interval

By substitution:

$$n = \frac{170}{1+170(0.05)^2} \quad n= 120$$

Hence, the sample size for the study = 120 respondents.

Using an actual population of 170 nurses, a sample size of 120 respondents were required for this study. Considering a 10% non-response rate, an estimated sample size of 132 was considered sufficient in powering the present study. Therefore, a total of 132 nurses were invited into the study.

2.5.1 Data collection instrument

A structured questionnaire was used for data collection with Pediatric Nurses' Knowledge and Attitudes Survey Regarding Pain protocol in focus. Information on participants socio-demographic data, nurses' knowledge and attitude regarding children's pain management were assessed.

2.5.2 Data collection procedure

After obtaining the requisite approvals from the institution, the researchers approached nurses at their recess time between duties at the hospital

under study to brief them about the study. The aims and nature of the study was explained to the participants before obtaining their informed consent. Those who consented to participate were given hard copies of the questionnaire which took about 20–30 minutes to complete over a one-month period in the year 2020.

2.6 Data Management

All data was entered into encrypted Microsoft Excel spreadsheets. To ensure confidentiality, only unique identifiers were included with the survey data in the electronic database. Encryption and passwords were used on the researchers' laptop. Files were deleted from the researchers' computer after study completion and were transferred to an external electronic drive, with the participant log and the survey data stored separately. The questionnaires were stored by the researchers in a locked filing cabinet at the Department of Paediatric Nursing and Child Health, University for Development Studies - Tamale.

2.7 Data Processing and Analysis

The IBM Statistics Package for Social Sciences (SPSS) Version 25 was used to analyze data. Participants' data was initially entered and cleaned in Microsoft excel before being exported into Statistical Package for Social Sciences version 25 for further analysis. Descriptive statistics, such as frequencies, percentages, means, and standard deviations used to describe and summarize the data. The Pearson Chi square test was used to determine the association between variables. A $p < 0.05$ was considered to be statistically significant. A correctly answered question on the Paediatric Nurses' Knowledge and Attitudes Survey Regarding Pain (PNKAS) instrument was awarded one point, whereas an incorrectly answered item will be given a zero score. The total score of this 42-item instrument will be converted into percentages for standardization purposes. A nurse who obtains 80% of the total score was considered as having satisfactory knowledge regarding children's pain.

3. FINDINGS

A total of 125 questionnaires were distributed over a 1-month period, 123 were returned, out of which 120 were valid and completed to be included in the analysis indicating a response rate of 96%. Sixty-two (62) representing fifty-seven percentage (51.7%) of participants were below 30 years of age and 58 (48.3%) of the

participants were aged 30 years and above. The majority 64 (53.3%) of participants were women and 56 (46.7%) were men. More than half of the participants 68 (56.7%) held diploma, and about one-fourth of participants 50(41.7%) had a bachelor's degree. Only 2(1.7%) nurses had a master's degree. The majority of participants 75 (62.5%) had less than 5 years of working experience in the Tamale West Hospital. A vast majority 110 (91.7%) of nurses reported not attending any pain management training/workshop. Most of the nurses were currently working in the medical wards 50 (41.7%). Only 20 (16.7%) of the nurses were currently working in pediatric ward. Most of the nurses 108 (90%) stated that there was no pain management standard or protocol used as a basis in the hospital. On the other hand, 12 (10%) stated that there was a pain management standard or protocol in the hospital. For the majority of nurses' their current position was senior staff nurse 85 (70.8%), and the rest of them were staff nurses 15 (12.5%), nursing officers 10 (8.3%), and senior nursing officers 10(8.3%), respectively.

3.1 Nurses Knowledge Regarding Children's Pain Management

As shown on Table 2, the top 10 items most often answered correctly by the participants. These items are centered on: the subjective and multidimensional nature of the pain experience and its treatment, the benefits of pre-emptive and multimodal analgesia, the role of parental presence during painful procedures, pain perceptions, useful pain medications and drug interactions.

4. DISCUSSION

4.1 Knowledge and Attitude on Pain Management

The current study aimed at assessing nurses' knowledge and attitudes regarding children's pain at the Tamale West Hospital in Ghana. Additionally, the researchers were interested in examining the socio-demographic factors that influence their pediatric pain knowledge and attitudes. The overall aim of this study was to provide baseline information to guide the development and implementation of a pediatric pain educational program for nurses to improve pediatric pain assessment and management. The study informed national and international literature by bringing new data about nurses in an area of limited knowledge.

Table 1. Results revealed that the mean score

Item contents	Mean (SD)
Comparable stimuli in different people produce the same intensity of pain (<i>F</i>)	3.94 (0.94)
Children who will require repeated painful procedures should receive maximum treatment for the pain and anxiety of the first procedure to minimize the development of anticipatory anxiety before subsequent procedures (<i>T</i>)	3.88 (0.97)
Combining analgesics and non-drug therapies that work by different mechanisms may result in better pain control with fewer side effects than using a single analgesic agent (<i>T</i>)	3.78 (0.99)
After the initial recommended dose of opioid analgesic, subsequent doses should be adjusted in accordance with the individual patient's response (<i>T</i>)	3.06(1.20)
Parents should not be present during painful procedures (<i>F</i>)	3.67(1.00)
Spiritual beliefs may lead a child/adolescent to think that pain and suffering are necessary (<i>T</i>)	3.63 (1.11)
Ibuprofen and other non-steroidal anti-inflammatory agents are NOT effective analgesics for bone pain caused by metastases (<i>F</i>)	3.60 (1.03)
Because their nervous system is underdeveloped, children under 2 years of age have decreased pain sensitivity and limited memory of painful experiences (<i>F</i>)	3.56 (1.04)
Pediatric patients who can be distracted from pain this usually do not have severe pain (<i>F</i>)	3.50(1.02)
Benzodiazepines do not reliably potentiate the analgesia of opioids' unless the pain is related to muscle spasms (<i>T</i>)	3.49 (1.04)
Non-drug interventions are very effective for mild-moderate pain control but are rarely helpful for severe pain (<i>F</i>)	3.40(1.14)
Giving children sterile water by injection (placebo) is often a useful test to determine if the pain is real (<i>F</i>)	3.37(1.35)
An equivalent of 15 mg of oral morphine is: (<i>Morphine 5 mg IV</i>)	3.30(1.23)
Observable changes in vital signs must be relied upon to verify a child's statement that he/she has severe pain (<i>F</i>)	3.31(1.11)
The percentage of patients who over report the amount of pain they have is: (0 or 10)	3.25(1.15)
Increasing analgesic requirements are signs that the patient is becoming addicted to the narcotic	3.09 (1.19)
About 25% of patients receiving narcotics around the clock become addicted	2.68(1.11)
Estimation of pain by a nurse of physician is a valid measure of pain as patients self-report	2.38(1.03)
The likelihood of narcotic addiction in children who are being treated with opioids is: (<1%)	2.36(0.98)
Which of the following drugs are potentially useful for treatment of children's cancer pain? (<i>All of the above</i>)	2.35(1.04)
Overall	3.19 (1.05)

Table 2. Top 10 items most often answered correctly by participants (n=120)

Item contents (correct answer)	f (%) Correct
Comparable stimuli in different people produce the same intensity of pain (<i>F</i>)	94 (78.3)
Children who will require repeated painful procedures should receive maximum treatment for the pain and anxiety of the first procedure to minimize the development of anticipatory anxiety before subsequent procedures (<i>T</i>)	93 (77.5)
Combining analgesics and non-drug therapies that work by different mechanisms may result in better pain control with fewer side effects than using a single analgesic agent (<i>T</i>)	84 (70)
After the initial recommended dose of opioid analgesic, subsequent doses should be adjusted in accordance with the individual patient's response (<i>T</i>)	83 (69.2)
Parents should not be present during painful procedures (<i>F</i>)	81 (67.5)
Spiritual beliefs may lead a child/adolescent to think that pain and suffering are necessary (<i>T</i>)	69 (57.5)
Ibuprofen and other non-steroidal anti-inflammatory agents are NOT effective analgesics for bone pain caused by metastases (<i>F</i>)	68 (56.7)
Because their nervous system is underdeveloped, children under 2 years of age have decreased pain sensitivity and limited memory of painful experiences (<i>F</i>)	67 (55.8)
Pediatric patients who can be distracted from pain this usually do not have severe pain (<i>F</i>)	67 (55.8)
Benzodiazepines do not reliably potentiate the analgesia of opioids' unless the pain is related to muscle spasms (<i>T</i>)	66 (55.0)

Note: f – Frequency; % – Percentage; T – True; F – False

The mean score of all responses was 3.12, indicating that participants' knowledge and attitudes were poor, corroborating findings from previous studies and reiterating the global concern about this issue among nurses who care for patients experiencing pain [16,17]. There appears to be an internationally recognized knowledge and attitudes deficits among nurses on this subject [18,19,20]. One of the possible reasons for this trend may be related to the insufficient pediatric pain management training in nursing curricula [21,22]. Limited continuing educational opportunities for nurses on this subject may also be responsible for these outcomes [23,24]. Due to these insufficiencies in knowledge and attitudes, pain in children globally remains inadequately managed, which leads to unnecessary suffering in this vulnerable population. The universal nature of the problem is reflected specifically in this study by the mean score achieved by nurses (3.12 in 63.9% of Jordanian nurses), which is close to that reported for nurses in the United Kingdom (3.20 in 65.7% of nurses in the United Kingdom) [25]. Other studies also reported generally poor nurses' knowledge of PM, where the mean scores fell in the range from 39.7% to 72.3% (Kassa & Kassa, 2014) [26,27]. There is an apparent reported deficit in nursing knowledge of pain management (PM) among nurses from different countries around the globe, and this is alarming. These mean scores mean that when nurses yield a score less than the threshold of 80%, their ability to care for patients experiencing pain might be considerably compromised [28,25], which is likely to contribute to unpleasant or negative patient experiences of PM, leading to low satisfaction with care.

A major concern observed in the current study was the inadequacy of knowledge and attitudes related to pain assessment, analgesics and pharmacokinetics (drug administration, distribution, metabolism and elimination), similar to previous studies [29,30,31]. Greater emphasis should be placed on these areas during the development and implementation of educational interventions on pediatric pain. Consistent with earlier studies [32,33,29] majority of the studied participants believed that the giving of placebo injection is a useful test for determining the veracity of patient's pain and equally perceived non-pharmacological interventions to be ineffective in managing severe pain. This suggests that the nurses did not believe in patient's self-report of pain despite the evidence that it is the best way of assessing and

evaluating a patient's pain [34]. It also further demonstrates that, majority of nursing and midwifery students harbour misconceptions about the efficacy on non-pharmacological pain management interventions. Non-pharmacological interventions have been shown to reduce pain associated with diverse causes and procedures [35]. The use of non-pharmacological interventions should therefore be encouraged as they reduce the amount of analgesic consumption needed in treating pain and associated side effects [36,37,38]. In the present study, most of the nurses answered the questions about pharmacology and addiction incorrectly. This finding corroborates a study by Alotaibi et al. [2] where the majority of nurses had insufficient knowledge regarding pharmacological management of pain. Education on pharmacological management of pain needs to be intensified in nursing curricula and regular in-service training organized for practicing nurses to enhance their knowledge on pharmacological interventions in pain management.

Consistent with a study by Amponsah et al. (2020), the present study found out that all the participants could not accurately determine the onset of action of orally administered analgesics, equianalgesic dose of orally administered morphine, and the right dosage of prescribed morphine for a child who consistently reported of moderate to severe pain. These results are of great concern as these nurses are already practicing in hospitals and may end up mismanaging children's pain. As the problem of poor assessment and management of children's pain persists among nurses, there is an urgent need to develop, implement and evaluate pain educational interventions especially in resource constrained setting to improve pain management for vulnerable children and their families.

4.1.1 Association between participants' characteristics and pain management

Although the mean scores of nurses in knowledge and attitudes scales were poor overall, further analysis revealed some statistical differences among different personal characteristics. Participants aged more than 30 years scored higher on the attitudes scale ($P5 .001$), similar to findings of studies conducted in Zimbabwe and Malaysia, where nurses aged greater than 40 years had better knowledge of PM. [26] (Ho, Ho & Pang, 2015).

There was also an association between years of experience and knowledge about PM, which is

supported by earlier research findings [39] (Lui, So & Fong, 2008). In contrast, a study in Turkey [39] found a negative correlation between the mean scores on the attitudes scale, and both age and working experience as older nurses had lesser positive attitudes toward PM, and less years of experience were also associated with lower mean scores on the scale. In this context, this might indicate that older nurses are more likely to hold negative attitudes and poor knowledge regarding PM. Nevertheless, other studies revealed that there was no significant difference among nurses' pain knowledge among subgroups for variables such as nurses' age, nursing experience years, and ranks (Lobo & Martins, 2013). In the present study, older and more experienced nurses scored higher compared with their younger counterparts, which may be related to the post qualification education within the target hospital, as senior nurses with adequate training in PM are often the decision makers about pain relief on the wards. There were significant differences in participants' attitudes toward PM related to the academic degree in nursing, in favor of those who have master's degrees in nursing. However, the number of master holders in this study is too small (only 2 participants) to draw meaningful conclusions for clinical practice, but it certainly indicates that further advanced education positively changes or reshapes nurses' knowledge and attitudes toward PM. Again, it is too early to make such conclusions and further investigation into this issue is warranted. Indeed, hospital-based PM training has been found to be associated with better decision-making processes, knowledge, and attitudes, consequently promoting pain relief among patients [40,41].

Previous studies found that women were more sensitive to the experience of pain than men [42,43]. As in most cultures, women generally adopt the family caregiver role in northern Ghana, which might endow them with latent skills and folk knowledge in dealing with illness-associated suffering and pain. Indeed, unlike male nurses, female nurses in Ghana look after both male and female patients, and presumably this might maximize their knowledge base and skills in pain management. However, at any rate, the mean scores of male nurses indicate the need for additional assistance to improve the efficacy of their pain management role within the healthcare context.

5. CONCLUSION

The study revealed that nurses at the Tamale West Hospital in Ghana generally have insufficient knowledge and attitudes toward pain in children. There is the urgent need to intensify education in this area so as to adequately prepare these nurses to cater for the pain needs of vulnerable children and their families. Future studies should focus on how student's theoretical knowledge and attitudes are linked to their pediatric pain assessment and management practices.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT

The study protocol and participants information sheet were provided for participants to read. The aim and objectives of the study was explained to participants and written informed consent obtained prior to data collection. The participants were informed that participation was solely voluntary and no monetary compensation was given to respondents for participating in the study. Respondents were assured of anonymity and confidentiality of their responses. Respondents were informed that they can opt out of the study at any point without any penalty and their decision would not have any consequences on them. Respondents were assured of confidentiality and that no personal identifiers will be used in the questionnaire. The nurses were informed that the study does not pose any risk to them. No compensation or monetary reimbursement was given for participating in the study.

ETHICAL APPROVAL

Permission to gain access to nurses was secured from the management of the Tamale West Hospital.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Shoqirat N, Mahasneh D. Nurses' Knowledge, Attitudes, and Barriers Toward Pain Management Among Postoperative Patients in Jordan. *Journal of PeriAnesthesia Nursing*. 2019;34 (2):359–367.
Available:https://doi.org/10.1016/j.jopan.2018.05.012
2. Alotaibi K, Higgins I, Chan S. Pain Management Nursing Nurses' Knowledge and Attitude toward Pediatric Pain Management: A Cross-Sectional Study. *Pain Management Nursing*. 2019;20(2): 118–125.
Available:https://doi.org/10.1016/j.pmn.2018.09.001
3. Ramira ML, Instone S, Clark MJ. Quality improvement. *Pediatric pain management: An evidence-based approach*. *Pediatric Nursing*. 2016;42(1):39e49.
4. Lunsford L, Mogan. Original Article Knowledge and Attitudes Regarding Pediatric Pain in Mongolian Nurses. *Pain Management Nursing*. 2015;16(3):346–353.
Available:https://doi.org/10.1016/j.pmn.2014.08.007
5. Boateng KHT, Yakong VN, Yombeï LN. Exploring Parents Needs Among Newborn Babies at Tamale Teaching Hospital. *Asian Journal of Pregnancy and Childbirth*. 2021;4(4):68-75.
Available:https://www.journalajpcb.com/index.php/AJPCB/article/view/30152
6. Beltramini A, Milojevic K, Pateron D. Pain assessment in newborns, infants, and children. *Pediatr Ann*. 2017;46(10):e387–95.
Available:http://www.ncbi.nlm.nih.gov/pubmed/29019634
7. Gerrits MM, van Oppen P, Leone SS, van Marwijk HW, van der Horst HE, Penninx BW. Pain, not chronic disease, is associated with the recurrence of depressive and anxiety disorders. *BMC Psychiatry* [Internet]. 2014 [cited 2019 Sep 1];14(1):187.
Available:http://bmcp psychiatry.biomedcentral.com/articles/10.1186/1471-244X-14-187
8. Dueñas M, Ojeda B, Salazar A, Mico JA, Failde I. A review of chronic pain impact on patients, their social environment and the health care system. *Journal of Pain Research*. 2016;9:457–67.
DOI:10.2147/JPR.S105892.
9. Tennant F. The physiologic effects of pain on the endocrine system. *Pain Ther* [Internet]. 2013;2(2):75–86.
Available:http://www.ncbi.nlm.nih.gov/pubmed/25135146
10. Hasuo H, Kusunoki H, Kanbara K, Abe T, Yunoki N, Haruma K, et al. Tolerable pain reduces gastric fundal accommodation and gastric motility in healthy subjects: a crossover ultrasonographic study. *Biopsychosoc Med* [Internet]. 9] 2017;11(1):4.
Available:http://bpsmedicine.biomedcentral.com/articles/10.1186/s13030-017-0089-5
11. Tumin D, Drees D, Miller R, Wrona S, Hayes D, Tobias JD, et al. Health care utilization and costs associated with pediatric chronic pain. *J Pain* [Internet]. 2018 [cited 2019 Sep 1];19(9):973–82.
Available:http://www.ncbi.nlm.nih.gov/pubmed/29608973
12. Agoston A, Gray L, Logan D. Pain in school: patterns of pain-related school impairment among adolescents with primary pain conditions, juvenile idiopathic arthritis pain, and pain-free peers. *Children* [Internet]. 2016 [cited 2019 Sep 1];3(4): 39.
Available: http://www.mdpi.com/2227-9067/3/4/39
13. Olmstead DL, Scott SD, Austin WJ. Unresolved pain in children: a relational ethics perspective. *Nurs Ethics* [Internet]. 2010 [cited 2019 Sep 1];17(6):695–704.
Available:http://journals.sagepub.com/doi/10.1177/0969733010378932
14. Al-Khawaldeh OA, Al-Hussami M, Darawad M. Knowledge and attitudes regarding pain management among Jordanian nursing students. *Nurse Educ Today*. 2013;33(4):339–45.
15. Yamane T. *Statistics: An introductory analysis*. 3rd ed. New York, USA: Harper and Row;1967.
16. Gupta M, Sahi MS, Bhargava A, Talwar V. The prevalence and characteristics of pain in critically ill cancer patients: A prospective nonrandomized observational study. *Indian J Palliat Care*. 2015;21:262-266.
17. Schroeder DL, Hoffman LA, Fioravanti M, Poskus, Medley D, Zullo TG, Tuite PK. Enhancing nurses' pain assessment to

- improve patient satisfaction. *Orthopaedic Nurs.* 2016;35:108-112.
18. Smeland AH, Twycross A, Lundeberg S, Rustøen T. Nurses' knowledge, attitudes and clinical practice in pediatric postoperative pain management. *Pain Manag Nurs.* 2018;19:585–598. Available:<https://doi.org/10.1016/j.pmn.2018.04.006> PMID: 29778756
 19. Stanley M, Pollard D. Relationship between knowledge, attitudes, and self-efficacy of nurses in the management of pediatric pain. *Pediatr Nurs.* 2013;39:165–171. Available:<http://www.ncbi.nlm.nih.gov/pubmed/24027950> PMID: 24027950
 20. Ekim A, Ocakci AF. Knowledge and attitudes regarding pain management of pediatric nurses in turkey. *Pain Manag Nurs.* 2013;14:e262–e267. Available:<https://doi.org/10.1016/j.pmn.2012.02.004> PMID: 24315279
 21. Mackintosh-Franklin C. Pain: A content review of undergraduate pre-registration nurse education in the United Kingdom. *Nurse Educ Today.* 2017;48:84–89. Available:<https://doi.org/10.1016/j.nedt.2016.08.035> PMID: 27723538
 22. Twycross A, Roderique L. Review of pain content in three-year preregistration pediatric nursing courses in the United Kingdom. *Pain Manag Nurs.* 2013;14:247–258. Available:<https://doi.org/10.1016/j.pmn.2011.05.002> PMID: 24315248
 23. Samarkandi O. Knowledge and attitudes of nurses toward pain management. *Saudi J Anaesth.* 2018;12: 220. Available:https://doi.org/10.4103/sja.SJA_587_17 PMID: 29628831
 24. Eid T, Manias E, Bucknall T, Almazrooa A. Nurses' knowledge and attitudes regarding pain in Saudi Arabia. *Pain Manag Nurs.* 2014;15:e25–e36. Available:<https://doi.org/10.1016/j.pmn.2014.05.014> PMID:25179424
 25. Vickers N, Wright S, Staines A. Surgical nurses in teaching hospitals in Ireland: Understanding pain. *Br J Nurs.* 2014;23:924-929.
 26. A, Chipfuwa T, Mukwamba MM, Chironda G. Knowledge and attitudes of registered nurses towards pain management of adult medical patients: A case of Bindura Hospital. *Health Sci.* 2015;9:1-6.
 27. Omran S, Al Qadire M, Ali NA, Al Hayek MF. Knowledge and attitudes about pain management: A comparison of oncology and non-oncology Jordanian nurses. *Nurs Health.* 2014;2:73-80.
 28. McCaffery M, Ferrell BR, Pasero C. Nurses' personal opinions about patients' pain and their effect on recorded assessments and titration of opioid doses. *Pain Manag Nurs.* 2011;1(3):79–87. Available:<http://www.ncbi.nlm.nih.gov/pubmed/11706463>
 29. Ortiz MI, Ponce-Monter HA, Rangel-Flores E, Castro-Gamez B, Romero-Quezada LC, O'Brien JP, et al.. Nurses' and nursing students' knowledge and attitudes regarding pediatric pain. *Nurs Res Pract [Internet].* [cited 2019 May 22]. 2015(210860):1–8. Available:<http://www.ncbi.nlm.nih.gov/pubmed/26543643>
 30. Al Omari O. Knowledge and attitudes of Jordanian nursing students toward children's pain assessment and management: A cross-sectional study. *J Nurs Educ Pract.* 2016;6(3):51–8.
 31. Gadallah MAEA, Hassan AMH, Shargawy SAEH. Undergraduate nursing students' knowledge and attitude regarding pain management of children in Upper Egypt. *J Nurs Educ Pract [Internet].* 2017;7(6):100–7. Available:<http://www.sciedupress.com/journal/index.php/jnep/article/view/10557>
 32. Kheshti R, Namazi S, Mehrabi M, Firouzabadi D. Health care workers' knowledge, attitude, and practice about chronic pain management, Shiraz, Iran. *Anesthesiol Pain Med [Internet].* 2016 [cited 2019 Sep 1];6(4):e37270. Available:<http://www.ncbi.nlm.nih.gov/pubmed/27843775>
 33. Becker WC, Dorflinger L, Edmond SN, Islam L, Heapy AA, Fraenkel L. (Bernardes SF, Silva SA, Carvalho H, Costa M, Pereira S. Is it a (fe)male pain? Portuguese nurses' and laypeople's gendered representations of common pains. *Eur J Pain.* 2014;18:530-539.
 34. McCaffery M, Ferrell BR, Pasero C. Nurses' personal opinions about patients' pain and their effect on recorded assessments and titration of opioid doses. *Pain Manag Nurs [Internet].* 2000;1(3):79–87.

- Available:<http://www.ncbi.nlm.nih.gov/pubmed/11706463>
35. Tick H, Nielsen A, Pelletier KR, Bonakdar R, Simmons S, Glick R, et al. Evidence-based nonpharmacologic strategies for comprehensive pain care. *EXPLORE* [Internet]. 2018;14(3):177–211. Available:<http://www.ncbi.nlm.nih.gov/pubmed/29735382>
36. McSherry T, Atterbury M, Gartner S, Helmold E, Searles DM, Schulman C. Randomized, crossover study of immersive virtual reality to decrease opioid use during painful wound care procedures in adults. *J Burn Care Res*. 2019;39(2):278–85. Available:<https://academic.oup.com/jbcr/article/39/2/278-285/4796748>
37. Rosendahl J, Koranyi S, Jacob D, Zech N, Hansen E. Efficacy of therapeutic suggestions under general anesthesia: A systematic review and meta-analysis of randomized controlled trials. *BMC Anesthesiol* [Internet]. 2016 [cited 2019 Sep 1];16(1):125. Available:<http://bmcanesthesiol.biomedcentral.com/articles/10.1186/s12871-016-0292-0>
38. Grissa MH, Baccouche H, Boubaker H, Beltaief K, Bzeouich N, Fredj N, et al. Acupuncture vs intravenous morphine in the management of acute pain in the ED. *Am J Emerg Med* 2016;34(11):2112–6. Available:<https://linkinghub.elsevier.com/retrieve/pii/S0735675716304223>
39. Manwere A, Chipfuwa T, Mukwamba MM, Chironda G. Knowledge and attitudes of registered nurses towards pain management of adult medical patients: A case of Bindura Hospital. *Health Sci*. 2015;9:1-6.
40. Silva MA, Pimenta CA, Cruz Dde A. Pain assessment and training: The impact on pain control after cardiac surgery. *Rev Esc Enferm*. 2013;47:84-92. 50.
41. Albaqawi H, Maude P, Shawhan-Akl L. Saudi Arabian nurses' knowledge and attitudes regarding pain management: Survey results using the KASRP. *Int J Health Sci Res*. 2016;6:150-164.
42. Bernardes SF, Silva SA, Carvalho H, Costa M, Pereira S. Is it a (fe)male pain? Portuguese nurses' and laypeople's gendered representations of common pains. *Eur J Pain*. 2014;18: 530-539.
43. Koren L, Peled E, Trogan R, Norman D, Berkovich Y, Israelit S. Gender, age and ethnicity influence on pain levels and analgesic use in the acute whiplash injury. *Eur J Trauma Emerg Surg*. 2015;41:287-291.

© 2021 Yombeï et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/78229>