



# Traumatic Extradural Hematoma and Kernohan's Phenomena in Pediatric Patient: Case Report and Current Concepts

Farah Wahidah Sazali <sup>a,b,c\*</sup> and Mohamad Azhari Bin Omar <sup>c</sup>

<sup>a</sup> Department of General Surgery, Universiti Kebangsaan Malaysia, Malaysia.

<sup>b</sup> Department of General Surgery, Hospital Selayang, Malaysia.

<sup>c</sup> Department of Neurosurgery, Hospital Raja Permaisuri Bainun, Ipoh, Malaysia.

## **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

## **Article Information**

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114024>

**Case Report**

**Received: 25/12/2023**

**Accepted: 27/02/2024**

**Published: 01/03/2024**

## **ABSTRACT**

**Introduction:** Traumatic extradural hemorrhage is one of the types of brain injury in children. The mortality of this type of brain injury can be as high 11.9% where the fall is the most common mechanism. Immediate surgical intervention such as decompressive craniectomy is a gold standard treatment in extradural hemorrhage.

**Case Report:** This is a case report of pediatric patient in trauma that sustained extradural hemorrhage that require surgical intervention due to low initial GCS upon presentation. Post trauma patient sustained ipsilateral weakness which was late recognition due to rare complication.

**Conclusion:** Managing extradural hemorrhage in pediatric patient is quite challenging as it has diversity of presentations where immediate surgical intervention is needed to prevent further complications.

\*Corresponding author: E-mail: farahwahidahsazali@gmail.com;

**Keywords:** Kernohan's phenomena; KWNP; extradural hematoma; traumatic brain injury pediatric head trauma.

## 1. INTRODUCTION

Traumatic Head Injury in pediatric population in Malaysia is about 32 per 100,000 children. A study done in Tay et al in 2016 showed pediatric head injury rate in Emergency Department admission accounted for 12.7% [1]. From this data, 11.9% was reported as mortality rate. Among children aged 9 and under, falls were the predominant cause of traumatic brain injuries [1]. Delay in surgical intervention in pediatric age group can lead to mortality and serious morbidity such as ischemia of brain and ipsilateral hemiparesis [2]. This case report focused on Traumatic Extradural Hemorrhage (EDH) in pediatric group with sequelae.

## 2. CASE PRESENTATION


A four-year-old child with a history of an alleged fall while being carried by her brother on his shoulder accidentally fell onto the floor from a height of approximately one meter. Following the trauma, the child experienced vomiting but did not seek medical attention. Five hours after the incident, she was found to be less responsive and in a tonic position over her right upper and lower limbs. Upon initial presentation to the emergency department, her Glasgow Coma Scale (GCS) was poor [3], and she had unequal pupils, resulting in intubation for airway protection. The primary and secondary surveys revealed no significant injuries other than a large temporoparietal epidural hematoma measuring 3

cm thickness with midline shift; therefore, she underwent left decompressive craniectomy and clot evacuation.


Intraoperative findings showed left temporoparietal EDH thickness 3cm evacuated. Repeated CT scans of the brain post operatively revealed acute infarcts in both left occipital lobe and left anteromedial temporal lobe. Furthermore, post-operative imaging indicated focal hypodensity involving the right crus cerebri due to mass effect from pre-operative CT scan results. On day six following surgery, child manage to wean off sedation and it was discovered that the child had developed left hemiparesis with upper limb hypotonia and hypertonia in her left lower limb. She successfully extubated on day ten with full GCS recovery but still exhibited residual left hemiparesis at discharge which required management by rehabilitation specialists.

## 3. DISCUSSION

Traumatic Brain Injury in pediatric population can result in morbidity and mortality. The annual mortality rate for children younger than 4 years old approximately 5 per 100000 and is higher compare to those in 5-14 years of age [3]. However the mortality rate between these pediatric age group can be as high 5.8% to 11.9% [1,4]. Most of the study showed that male is the predominance group in pediatric population compare to female [3-5].



**Fig. 1. a. Initial CT scan showed large temporoparietal EDH. b. Size of EDH showed 3 cm with midline shift 0.8cm**



**Fig. 2. Showed post operative CT scan. On right side noted focal area of hypodensity in right crus cerebri in midbrain. On left sided showed hypodensities over left anteromedial temporal lobe and occipital lobe representing infarcts**

Extradural hematoma (EDH) is one of type of brain injury where there is an accumulation of blood between inner table of skull and the stripped off dural membrane consist of venous blood [6]. Majority of traumatic venous EDH manifest in pediatric patients are typically not correlated with any accompanying skull fractures [7]. In infants group, EDH is rare because the dura mater is adhered strongly to inner surface of skull especially in proximity suture line [3].

The study conducted by Browne and Lam found that 45.5% of patients had isolated EDH with no associated injury and fall as a major factor. A new study done by Mamoori et al in Pakistan showed almost similar results with 38.8% of cases being isolated EDH, with falls contributing to half of these cases [7,8]. For the evaluation of EDH, plain skull x-ray is not helpful. CT scan is the gold standard with appearance of high density biconvex shape can be seen in 84% of the cases. Surgery is indicated when the EDH is more than 30cc regardless of GCS. Non-surgical intervention may be preferred for cases that exhibit a volume less than 30cc, thickness under 15mm, midline shift below 5mm and GCS score more than eight in the absence of focal deficits and should be monitored in neurosurgical centre. In the TBI pediatric population, individuals presenting to the ED with a reduced GCS accounted for approximately 56.6% of mortality rates within a year. Nonetheless, an additional 15% of patients experienced good prognosis

over a period spanning at least ten years [3].

In this case study, the child successfully prevented mortality yet she still experienced morbidity characterized by left-sided hemiparesis consistent with the location of the bleeding side. This sign can be attributed to false localizing sign or Kernohan's notch Phenomena. Kernohan's phenomena was first described by James Kernohan in 1929 while writing the article known as incisura of the crus due to contralateral brain tumor [9]. This phenomena refers to an intracranial anomaly that results in a significant lateral mass effect, resulting in the compression of the contralateral cerebral peduncle against the free margin of the cerebellar tentorium [10]. It is also resulting from the compression of the contralateral cerebral peduncles against the tough dural reflections, which causes damage to pyramidal tract fibers located in the brainstem [11]. This ultimately leads to paralysis on the ipsilateral side of a lesion present in the brain as observed in our patient in this case report.

The most credible pathophysiological theory for KWNP suggests a transient inadequacy in regional blood circulation due to mechanical compression at the tentorial notch. In the study done by Beuclear et al in 2022 showed that 53% patients suffered from post operative neurological status whereby 29% of them

sustained ipsilateral hemiparesis and 18% had ipsilateral hemiplegia [10]. As for abnormality in pupil examination 8% suffered from ipsilateral mydriasis on arrival however were resolved later. It was reported that 92% were benefit from MRI study to diagnose Kernohan's phenomena [10]. Until recently, computed tomography (CT) scans were not utilized for diagnosing Kernohan's notch. However, when obtained to diagnose an intracranial space-occupying lesion, a CT scan can detect contralateral midbrain compression and, combined with clinical features indicating ipsilateral pyramidal tract involvement, provide support for the diagnosis of Kernohan's notch [11]. A solitary case study conducted by Ragesh et al. demonstrates that the efficacy of CT scans is sufficient, rendering additional MRI procedures unnecessary [11]. This findings also showed that CT scan also can be one of the modality of imaging in which centers that do not provide MRI scans. Based on study done by Beucler et al, 45% of patients achieved positive outcomes while 39% experienced negative outcomes [10]. Within the latter group, it was reported that an unfortunate of 22% suffered from hemiparesis. As for long term functional outcome statistically not associated with any factors such as age, dilated pupil upon admission or trauma lesion. The sole variable that lacked significance, yet exhibited a propensity towards improved motor functionality was a GCS score more than 3 upon admission [10].

#### 4. CONCLUSION

Managing traumatic EDH in pediatric patients can be a complex task due to the diverse clinical presentations that are age-dependent. Early recognition of this neurosurgical emergency is crucial to prevent mortality and minimize morbidity. Therefore, meticulous clinical evaluation and radiology assessments play a pivotal role in detecting early signs of morbidity.

#### CONSENT

As per international standards, parental written consent has been collected and preserved by the author(s).

#### ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Tay EL, Lee SWH, Jamaluddin SF, Tam CL, Wong CP. The epidemiology of childhood brain injury in the state of Selangor and Federal Territory of Kuala Lumpur, Malaysia. *BMC Pediatrics*. 2016; 16(1):56.
2. Ali MFA, Elbaroody M, Alsawy MF, El Fiki A, El Refaae E, Elshitany HA. Postoperative cerebral infarction after evacuation of traumatic epidural hematoma in children younger than two years: Single-center experience. *Surgical Neurology International*. 2022;13.
3. Araki T, Yokota H, Morita A. Pediatric Traumatic Brain Injury: Characteristic Features, Diagnosis, and Management. *Neurol Med Chir (Tokyo)*. 2017;57(2):82-93.
4. Hasanein M. Pediatric Extradural Hematoma: A Report of 34 Cases. *Suez Canal University Medical Journal*. 2019;22: 46-55.
5. Jamjoom A, Cummins B, Jamjoom ZA. Clinical characteristics of traumatic extradural hematoma: A comparison between children and adults. *Neurosurg Rev*. 1994;17(4):277-81.
6. Umerani MS, Abbas A, Aziz F, Shahid R, Ali F, Rizvi RK. Pediatric Extradural Hematoma: Clinical Assessment Using King's Outcome Scale for Childhood Head Injury. *Asian J Neurosurg*. 2018;13(3): 681-4.
7. Al-Mamoori M. Management of epidural hematoma in the pediatric age group. *Medical Journal of Babylon*. 2019;16:276.
8. Spazzapan P, Krašovec K, Velnar T. Risk factors for bad outcome in pediatric epidural hematomas: A systemic review. *Chin Neurosurg J*. 2019;5:19.
9. Etienne D, Zurada A, Matusz P, Shoja MM, Tubbs RS, Loukas M. James Watson Kernohan (1896–1981): Frontiers in neuropathology. *Clinical Anatomy*. 2012; 25(4):527-9.
10. Beucler N, Cungi PJ, Baucher G, Coze S, Dagain A, Roche PH. The Kernohan-Woltman Notch Phenomenon: A Systematic Review of Clinical and Radiologic Presentation, Surgical

- Management, and Functional Prognosis. J Korean Neurosurg Soc. 2022;65(5):652-64.
11. Panikkath R, Panikkath D, Lim SY, Nugent K. Kernohan's Notch: A Forgotten Cause of Hemiplegia-CT Scans Are Useful in This Diagnosis. Case Rep Med. 2013;2013:296874.

---

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/114024>