



# Association between Stress Levels and Functional Dyspepsia among Medical Students During the COVID-19 Pandemic: A Study of Indonesian Christian University

Nur Nunu Prihantini <sup>a\*</sup>, Elshara Gabriela Kuheba <sup>a</sup>  
and Mulyadi Djojoputro <sup>a</sup>

<sup>a</sup> Faculty of Medicine, Universitas Kristen Indonesia, Jakarta, Indonesia.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijtdh/2024/v45i101597>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119032>

**Original Research Article**

**Received: 01/08/2024**

**Accepted: 03/10/2024**

**Published: 07/10/2024**

## ABSTRACT

**Background:** Stress is a problem that often occurs among students. The high incidence of stress in students can interfere with students' effectiveness in carrying out their lecture activities. Stress in students not only affects the quality of learning but can also affect the physical condition of students, one of which is the occurrence of gastrointestinal disorders, one example of which is dyspepsia.

\*Corresponding author: Email: [nur.nunu@uki.ac.id](mailto:nur.nunu@uki.ac.id);

**Cite as:** Prihantini, Nur Nunu, Elshara Gabriela Kuheba, and Mulyadi Djojoputro. 2024. "Association Between Stress Levels and Functional Dyspepsia Among Medical Students During the COVID-19 Pandemic: A Study of Indonesian Christian University". *International Journal of TROPICAL DISEASE & Health* 45 (10):46-54. <https://doi.org/10.9734/ijtdh/2024/v45i101597>.

**Aims:** This study aims to determine whether there is a relationship between stress levels and functional dyspepsia in students at the Faculty of Medicine, Indonesian Christian University class of 2020.

**Study Design:** This research uses an observational analytical research design method with a cross-sectional approach. The main data source is primary data obtained through questionnaires taken online via Google Forms.

**Place and Duration of Study:** The research was conducted at the Indonesian Christian University located in Cawang, East Jakarta. Time of research in September 2022.

**Methodology:** The population in this study was 137 students from the Faculty of Medicine, Indonesian Christian University class of 2020. The sample was taken using a simple random sampling technique totaling 103 people determined using the Slovin formula with an uncertainty percentage of 5%.

**Results:** From the results of data collection on 103 respondents who were FK UKI students Class of 2020, around 65% or 67 respondents were identified as having normal stress levels and as many as 64 students (62%) had symptoms of functional dyspepsia. Based on the results of the bivariate analysis, it was found that there was a relationship between the level of stress and the incidence of dyspepsia experienced by FK UKI students class of 2020 with the results of the chi-square test, namely  $p = 0.003 < \text{sig } 0.05$

**Conclusion:** There is a significant relationship between stress levels and functional dyspepsia. This certainly cannot be taken lightly because the incidence of functional dyspepsia can have fatal consequences which can disrupt the sufferer's activities and productivity. Therefore, it is very necessary to manage emotions (stress levels) both by the sufferer himself and with professional help.

*Keywords: Functional dyspepsia; stress levels; incidence of dyspepsia.*

## 1. INTRODUCTION

Stress is a condition characterized by the appearance of physical or emotional symptoms as a reaction to a person's feelings towards an undesirable situation [1]. Low levels of stress can be beneficial and can even improve bodily health. Stress can positively improve a person's biopsychosocial health and help improve performance. In addition, stress can act as a stimulus for self-development. However, if stress levels become too high, it can cause biological, psychological, and social problems, and even pose serious threats [2]. Stress can be triggered by stressors. Some examples of stressors include work, education, financial and family problems [3]. Stress has become a problem that exists at all levels of society. When someone experiences excessive stress, it can harm their physical condition, one of which is affecting the gastrointestinal system which can cause dyspepsia. Dyspepsia is a term that comes from Greek, where "dys" means bad and "pepsis" refers to digestion [4]. It is a collection of symptoms that include a burning sensation in the upper part of the stomach, pain, bloating, and nausea, vomiting [5]. Dyspepsia is divided into two types based on what the cause is. Functional Dyspepsia falls into the category of dyspepsia whose cause is unknown no abnormalities were

found on conventional gastroenterological examination or no organic damage was found [5]. Globally, the prevalence of dyspepsia is between 10-30% of the world's total population. The prevalence rate of dyspepsia in Indonesia reaches 40-50% and has increased significantly from 10 million people in 2020 to 28 million people, or around 11.3% of Indonesia's total population [6]. Data from Profile Indonesian Health in 2020 shows that dyspepsia has reached the 10th position as the digestive disease most frequently experienced by inpatients in hospitals in 2019, with the number of patients around 34,039 or around 1.59% of the total patients [7]. Students as part of society also experience stress, especially currently during a pandemic which causes learning to be carried out online. The stress level for students in Indonesia during distance lectures reached 55.1% while for students outside Indonesia, it reached 66.3% [8]. The high level of stress also contributes to the large number of cases of gastritis in Indonesia. This is what motivated the author to further research the relationship between stress levels and the incidence of gastritis in FK UKI Class of 2020 students [9].

**Research problem:** The research problem is whether there is a relationship between stress levels and functional dyspepsia in students from

the Faculty of Medicine, Indonesian Christian University class of 2020.

**Research purpose:** The research aims to find out whether there is a relationship between stress levels and functional dyspepsia in students at the Faculty of Medicine, Indonesian Christian University, class of 2020.

## 2. MATERIALS AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

### 2.1 materials

**Definition of stress:** Stress has many different definitions depending on the condition. One of the most common definitions of stress introduced by Hans Selye is that it is a non-specific response of the body to pressure received.<sup>9</sup> He also wrote that stress is the perception of threat, which produces worry, discomfort, emotional tension, and difficulty in adjustment [10]. According to McGrath (2003) Stress is an imbalance between physical or psychological demands and the ability to respond in conditions where: failure to fulfill those demands. This means that stress can arise in individuals if there is an imbalance or failure to meet physical and mental needs [11-14].

However, not everyone who experiences an imbalance will experience stress. This imbalance stimulus will be responded to differently in each individual. This happens because there is a difference in the ability to reduce the stimulus. Apart from that, there are also life experiences, the level of sensitivity, and the individual's tolerance for stimuli which also have an influence. So this makes each person's stress threshold different [15,16].

**Stress rating scale:** In this research, evaluation of stress levels was carried out through the use of an instrument in the form of a questionnaire. The measuring instrument used in this research is the Depression Anxiety Stress Scales (DASS). DASS or Depression Anxiety Stress Scales is a questionnaire to measure a person's symptoms of depression, anxiety and stress. This measuring instrument was developed by Lovibond in 1995 [17]. DASS has 3 measurement scales that support 13 items, which are divided into 2 to 5 subscales. The depression scale is used to evaluate mood which includes dysphoria, hopelessness, loss of interest in activities usually enjoyed, and

helplessness. Meanwhile, the anxiety scale is used to evaluate physical arousal, panic attacks, and fear. The stress scale consists of items that can evaluate the level of tension, irritability, and excessive reactions to certain events [18]. Instrument The DASS consists of 42 questions, and each question has a score ranging from 0 to 3. A score of 0 is given if the respondent never experiences certain symptoms, a score of 1 is given if they sometimes experience it, a score of 2 is given if they often experience it, and a score of 3 is given if it closely matches the respondent's experience or almost always occurs. Evaluation is carried out based on the respondent's condition for one week [19].

### Dyspepsia:

**Definition of dyspepsia:** Dyspepsia is a condition characterized by discomfort or pain in the upper part of the stomach or solar plexus [20] Dyspepsia refers to a group of signs and symptoms located in the epigastrium, where patients can experience heartburn or discomfort, a burning sensation, and may be accompanied by nausea, vomiting, bloating, early satiety, and belching. This collection of symptoms can appear sporadically over several weeks or even months. Not all patients have complaints, some patients only complain about one or more, and the type of complaint and quality can also change.

**Classification of dyspepsia:** Dyspepsia is categorized based on its cause and is divided into two parts, namely organic dyspepsia and functional dyspepsia [5,21].

#### a. Organic dyspepsia

Organic dyspepsia refers to dyspeptic conditions whose causes have been identified, such as peptic ulcer disease (PUD), gastric acid reflux disease (GERD), cancer, use of alcohol or drugs.

#### b. Non-organic dyspepsia

Non-organic dyspepsia, also called functional dyspepsia, is a condition in which dyspepsia occurs without any known structural or organic abnormalities as the cause. Functional dyspepsia is divided into two groups, namely Postprandial Distress Syndrome, which is a condition where a person feels full after eating a normal portion or is easily full so it is difficult to finish his meal. The second is Epigastric Pain Syndrome which refers to the condition of the patient experiencing a burning sensation. These symptoms come and go and do not occur in other parts of the stomach [22].

### 3. RESULTS AND DISCUSSION

#### 3.1 Results

##### 3.1.1 Respondent characteristics

The research respondents were students from the faculty of medicine, Indonesian Christian University, class of 2020, with active status, namely 137 people from the population as respondents in this research. Of the 137 students from the class of 2019, 21 people fell into the exclusion criteria so they could not continue to the research stage. So the respondents who filled out the questionnaire completely were 115 students, of whom 103 people were then selected as samples according to the simple random sampling method and the Slovin formula.

Based on univariate analysis, the frequency distribution of respondents based on age was obtained as presented in Table 1.

The information contained in the Table 1 shows that the majority of respondents, namely 52 people or around 50.3%, are 20 years old. Then followed by ages 21 years (25.2%), 19 years (22.3%), 18 years and 22 years, namely 1% for 1 person each.

The frequency distribution of respondents based on gender is presented in Table 2.

The gender distribution obtained from the 2020 FK UKI students based on the Table 2 is that female students numbered 79 respondents with

a percentage of 76.7% while male students totaled 24 respondents with a percentage of 23.2% so it can be concluded that the number of female students dominates more than men in FK UKI class of 2020. Meanwhile, the frequency distribution of respondents who experienced functional dyspepsia is presented in Table 3.

Based on the Table 3, it was found that most of the 2020 FK UKI students had functional dyspepsia where 64 respondents with a percentage of 63.1% had functional dyspepsia while 38 respondents with a percentage of 37.9% did not have functional dyspepsia

**Characteristics of respondents based on stress level:** The frequency distribution of respondents based on stress level is presented in Table 4.

The distribution of stress levels among FK UKI students class of 2020 is as follows: The number of students with normal stress levels, namely with a score of 0-14, is 67 respondents with a percentage of 65%. The number of respondents for mild stress levels was 12 people with a percentage of 11.7% followed by moderate stress levels with 16 people with a percentage of 15.5%. The lowest number of respondents was at the severe stress level and the very severe stress level, with 4 respondents each with a percentage of 3.9%. From the data obtained, it was concluded that the 2020 FK UKI students did not experience significant stress.

The distribution of stress levels based on gender is presented in Table 5.

**Table 1. The frequency distribution of respondents based on age**

Age	Frequency	Percentage (%)
18 years	1	1.0
19 years	23	22.3
20 years	52	50.5
21 years	26	25.2
22 years	1	1.0
<b>Total</b>	<b>103</b>	<b>100</b>

**Table 2. Frequency distribution based on gender**

Gender	Frequency	Percentage (%)
Male	24	23.3
Female	79	76.7
<b>Total</b>	<b>103</b>	<b>100</b>

**Table 3. Frequency distribution of respondents who experienced functional dyspepsia**

Functional Dyspepsia	Frequency	Percentage (%)
Experiencing Dyspepsia	64	62.1
Not Experiencing Dyspepsia	39	37.9
<b>Total</b>	<b>103</b>	<b>100</b>

**Table 4. Frequency distribution of respondents based on stress level**

Stress Level	Frequency	Percentage (%)
Normal (0-14)	67	65.0
Mild Stress (15-18)	12	11.7
Moderate Stress (19-25)	16	15.5
Severe Stress (26-33)	4	3.9
Very Severe Stress (>34)	4	3.9
<b>Total</b>	<b>103</b>	<b>100</b>

**Table 5. Frequency distribution of respondents regarding stress levels based on gender**

Stress Level	Amount		Total	Percentage (%)
	Male	Female		
<b>Normal</b>	20	47	67	65
<b>Mild Stress</b>	2	10	12	11.7
<b>Moderate Stress</b>	1	15	16	15.5
<b>Severe Stress</b>	1	3	4	3.9
<b>Very Severe Stress</b>	0	4	4	3,9
<b>Total</b>	<b>25</b>	<b>78</b>	<b>103</b>	<b>100</b>

**Table 6. Frequency distribution of stress levels based on age**

Level Stress	Age					Total	Perse ntase (%)
	18	19	20	21	22		
<b>Normal</b>	1	16	29	21	0	67	65
<b>Mild Stress</b>	0	3	5	3	1	12	11,6
<b>Moderate Stress</b>	0	1	13	2	0	16	15,5
<b>Severe Stress</b>	0	2	2	0	0	4	3,8
<b>Very Severe Stress</b>	0	1	3	0	0	4	3,8
<b>Total</b>	1	23	52	26	1	103	100

**Table 7. Distribution of functional dyspepsia based on age**

Age	Functional Dyspepsia		Total
	Ya	Tidak	
18 years	0	1	1
19 years	17	6	23
20 years	31	21	52
21 years	15	11	26
22 years	1	0	1
<b>Total</b>	<b>64</b>	<b>39</b>	<b>103</b>

**Table 8. The relationship between stress levels and functional dyspepsia**

Stress Level	Dyspepsia		Total
	Dyspepsia	Not Dyspepsia	
Normal	33	34	67
Mild Stress	10	2	12
Moderate Stress	14	2	16
Severe Stress	3	1	4
Very Severe Stress	4	0	4
<b>Total</b>	<b>64</b>	<b>39</b>	<b>103</b>
<b>P Value</b>		<b>0.003</b>	

The results of the study show that there are differences in the distribution of stress between genders, where women have a higher rate of experiencing stress. This is also supported by the fact that the population of FK UKI students class of 2020 is dominated by women. Based on the Table 5, the difference in stress levels between men and women is not very significant. Normal stress levels occupy the highest position, followed by moderate stress levels and light stress levels.

The frequency distribution of stress levels based on age is presented in Table 6.

Based on the Table 6, it can be concluded that the 20-year age group has the highest level of stress. Where the most is at the normal stress level, namely 29 out of 103 respondents in the 20-year age group.

Table 7. The age frequency distribution based on functional dyspepsia is presented in Table 7.

The distribution of functional dyspepsia based on age, namely in students aged 18 years, no dyspepsia was found. Then for 19-year-old students, of the 23 respondents, 17 respondents had dyspepsia, and 6 did not have dyspepsia. Continuing with students aged 20 years, 31 respondents had dyspepsia while 21 other respondents did not have dyspepsia. For students aged 21 years, 15 respondents had functional dyspepsia and 11 other respondents did not have dyspepsia. There was 1 respondent aged 22 years in students who had dyspepsia.

**Bivariate analysis:** The results of bivariate analysis to see the relationship between stress levels and functional dyspepsia are presented in Table 8:

Based on the Table 8, 33 respondents had normal stress levels and dyspepsia, and 31 respondents who did not have dyspepsia. There

were 10 respondents with mild levels of stress and dyspepsia and 2 respondents who did not have dyspepsia. The moderate level of stress with dyspepsia was 14 respondents and those without dyspepsia were 2 respondents. For the level of severe stress, 3 respondents had dyspepsia and 1 respondent did not have dyspepsia. All respondents with very severe stress levels had dyspepsia.

### 3.2 Discussion

**Univariate analysis:** Based on the results of univariate analysis research, it was found that 63% or 64 student respondents had functional dyspepsia. The emergence of functional dyspepsia cannot be viewed in isolation, because other factors in each individual also influence it. In theory, there are several factors such as diet (spicy food, drinks such as coffee, tea, and alcohol) and lifestyle such as smoking which are believed to contribute to the emergence of dyspepsia symptoms [6,23]. This theory is by research conducted by Clarensius Geovani and colleagues. in medical students at Nusa Cendana University, where there was a relationship between consuming spicy food, drinking tea or coffee, and carbonated drinks with the incidence of dyspepsia, namely with a p-value = 0.001. Other results were found in research on smoking habits and the incidence of dyspepsia syndrome conducted by Ivan Wijaya and colleagues, where the results found were a p-value = 0.190, indicating that there was no relationship between smoking habits and the incidence of dyspepsia. However, in the same study, it was found that there was a relationship between consuming alcoholic drinks and the incidence of dyspepsia with a value of  $p = 0.040$  [24].

The research results that were also obtained were that some students had normal stress levels, namely 65% with a total of 67 out of 103 respondents. This can be caused by differences

in each individual's ability to tolerate stress. The causes of stress in students can occur due to several wrong aspects the other is an academic problem. During the Covid-19 pandemic, students were required to carry out all lecture activities online, although online media had many shortcomings in its implementation. 6 According to research conducted by Alahan Durbas and his colleagues at 119 universities in Turkey, they found that there was a relationship between stress and anxiety with the Covid-19 pandemic. This research also found the causes of stress during the pandemic, namely due to socio-economic restrictions, online learning, and also concerns about the health of oneself and those closest to them [25].

**Bivariate analysis:** Based on the results of bivariate tests, there is a relationship between stress levels and the incidence of functional dyspepsia experienced by FK UKI Class of 2020 students. It was found that 31 students experienced mild to very severe levels of stress with functional dyspepsia while 31 students with mild to very severe levels of stress did not. dyspepsia as many as 33 people. Meanwhile, after analyzing the data using SPSS with the Chi-Square test, it was found that stress levels were related to functional dyspepsia. This can be seen from the results of the P value, namely  $p=0.003$ , which means  $p<0.05$  or the p value is smaller than the standard value. This value shows that the research results have a meaningful relationship. This research also proves that the results obtained by sisters Lady Maria and Salmah among students at Hasanuddin University in 2018 where the results of their research found a value of  $p = 0.019$  which means  $p < 0.05$  so the results of their research showed that there is a significant relationship. 44 Research also conducted by Prem Shankar and colleagues, it was found that students who experience stress have a risk of developing functional dyspepsia. The results of data analysis show a p-value = 0.001, where the p-value  $<0.005$  shows that the relationship between stress and dyspepsia has a significant correlation. The similarity in results in the incidence of functional dyspepsia can be influenced by various factors, such as the high level of stress experienced by many students, especially female students, the same age, and the same faculty is the medical faculty. Therefore, it cannot be concluded that the similarity of results is caused by one factor alone.

## 4. CONCLUSION

There is a significant relationship between stress levels and functional dyspepsia. This certainly cannot be taken lightly because the incidence of functional dyspepsia can have fatal consequences which can disrupt the sufferer's activities and productivity. Therefore, it is very necessary to manage emotions (stress levels) both by the sufferer himself and with professional help.

### Research limitations:

After conducting research, there were several limitations found by researchers. It is hoped that these findings can provide insight for future researchers to conduct better research. The following are some limitations found in this research:

1. Research questionnaires were not distributed actively (out in the field) because we were still in the COVID-19 pandemic, therefore, researchers could not directly observe the process of filling out the questionnaire.
2. Some respondents often answered with answers that were not in line with the previous statement. This can be caused by hasty and dishonest filling

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

## CONSENT AND ETHICAL APPROVAL

It is not applicable.

## ACKNOWLEDGEMENTS

We would like to express our gratitude to the Faculty of Medicine, Indonesian Christian University for the support and facilitation provided so that this research could run well.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Park CL, Russell BS, Fendrich M, Finkelstein-Fox L, Hutchison M, Becker J. Americans' COVID-19 stress, coping, and adherence to CDC guidelines. *J Gen Intern Med*. 2020, Aug 1;35(8):2296–303.
2. Madisch A, Andresen V, Enck P, Labenz J, Frieeling T, Schemann M. The diagnosis and treatment of functional dyspepsia. *Dtsch Arztebl Int*. 2018, Mar 30;115(13):222–32. databases Available:<http://www.ncbi.nlm.nih.gov/nlmcatalog/journals>.
3. Syahputra R, Siregar NP. The Relationship Between Stress Levels and the Incidence of Functional Dyspepsia Syndrome in Students of the Faculty of Medicine at the Islamic University of North Sumatra class of 2020. *Ibnu Nafis Medical Journal*. 2021, Dec 1;10(2):101–9. Available:<https://doi.org/10.30743/jkin.v10i2.178>
4. Sumarni S, Andriani D. The relationship of eating patterns with the incident of dyspepsia. *Journal of Nursing and Physiotherapy (JKF)*. 2019, Oct 31;2(1):61–6.
5. Fauziyyah R, Awinda RC, Besral B. The impact of distance learning on student stress and anxiety levels during the COVID-19 pandemic. *Journal of Biostatistics, Population and Health Informatics*. 2021, Mar 31;1(2):113–23.
6. Fink G. Stress: Concepts, Definition and History☆. In: *Reference Module in Neuroscience and Biobehavioral Psychology*. Melbourne: Elsevier. 2017;1–9.
7. Mofatteh M. Risk factors associated with stress, anxiety, and depression among university undergraduate students. *AIMS Public Health*. 2021;8(1):36–65. Available:<http://www.aimspress.com/article/doi/10.3934/publichealth.2021004>
8. Russell G, Lightman S. The human stress response. *Nat Rev Endocrinol*. 2019;15(9):525–34. Available:<https://doi.org/10.1038/s41574-019-0228-0>
9. Priyoto. *Stress management concept*. 2nd ed. Yogyakarta: Nuha Medika; 2019. 45–53 p
10. Wibawani EA, Faturahman Y, Purwanto A. Factors related to the incidence of dyspepsia in Internal Medicine Outpatient Clinic patients at Koja Regional Hospital. *Indonesian Community Health Journal*. 2021 Mar;17:258–64.
11. Mertz H. Stress and the Gut. UNC center for functional GI & motility disorders. 2017, Oct;1–5.
12. Shankar P, Mandhan N, Hussain Zaidi SM, Choudhry MS, Kumar A. Relationship of functional dyspepsia with mental and physical stress. *Annals of Psychophysiology*. 2020, Oct 1;7(1):25–30. Available:<https://aeirc-edu.com/ojs14/index.php/app/article/view/461>
13. Sugiono O. Validity and Reliability Test of the SG Posture Evaluation Measuring Instrument. *Journal of Physical Therapy*. 2020, May 4;5(1):55–61. Available:<https://jurnalketerapianfisik.com/index.php/jpt/article/view/167>
14. Wahyuningtyas A, Setiowati EA. Expressive writing to improve psychological well-being MA. *Psisula: Periodical Proceedings of Psychology*. 2020;2:17–33. Available:<https://journal.unnes.ac.id/nju/INTUISI/article/view/26388>
15. Lady M. The relationship between stress and eating habits on the incidence of dyspepsia among students at the Faculty of Medicine, Hasanuddin University Class of 2018. [Makassar]; 2018. Available:<https://e-journal.unair.ac.id/AMNT/article/view/45287>
16. Deani PS. THE Relationship of Eating Patterns and Stress Level to the Degree of Dyspepsia Syndrome In Adolescent Women At SMA N 3 Padang (Doctoral dissertation, Andalas University). Available:<http://scholar.unand.ac.id/69624/7/cover%20dan%20abstrak.pdf>
17. Muflih M, Najamuddin N. The relationship between diet and stress levels with the incidence of dyspepsia at Sundari Medan General Hospital in 2019. *Indonesian Trust Health Journal*. 2020, Dec 1;3(2):326-36. Available:<https://jurnal.murniteguhuniversit y.web.id/index.php/ithj/article/view/56>
18. Afifah N. The Relationship between Stress Levels and Diet with the Incidence of Dyspepsia Syndrome in Nursing Students in the Eighth Semester of 'Aisyiyah Yogyakarta University (Doctoral dissertation, 'Aisyiyah Yogyakarta University).



- Available:<https://digilib.unisayogya.ac.id/4393/1/naskah%20publikasi%20afifah-min.pdf>
19. Zulhikmah M. The Relationship of Characteristics, Eating Patterns, and Stress Level to the Incident Of Dyspepsi In The Pandawan Health Center, Hulu Sungai Tengah District; 2022. (Doctoral Dissertation, Kalimantan Islamic University, MAB). Available:<https://eprints.uniska-bjm.ac.id/12165/1/Artikel%20tya%20fix.pdf>
20. Dhita Da. The relationship of stress level to dyspepsia syndrome in faculty of medicine students, yarsi university, Class of 2018 and Its Review According to Islamic Views (Doctoral Dissertation, Yarsi University). Available:<https://digilib.yarsi.ac.id/11236/>
21. Maresa T. The Relationship of stress level and dietary regularity with the occurrency of dyspepsia in the productive age at the Depok Iii Puskesmas, Sleman Yogyakarta Card). All References Should Follow the Following Style.
- Available:[https://digilib.unisayogya.ac.id/4629/1/NASKAH%20PUBLIKASI\\_TRIA%20MARESA\\_1710201223.pdf](https://digilib.unisayogya.ac.id/4629/1/NASKAH%20PUBLIKASI_TRIA%20MARESA_1710201223.pdf)
22. Shankar P, Mandhan N, Hussain Zaidi SM, Choudhry MS, Kumar A. Relationship of functional dyspepsia with mental and physical stress. *Ann Psychophysiol.* 2020, Oct 1;7(1):25-30.
23. Huang ZP, Wang K, Duan YH, Yang G. Correlation between lifestyle and social factors in functional dyspepsia among college freshmen. *Journal of International Medical Research.* 2020, Aug;48(8):0300060520939702.
24. Gallas S, Knaz H, Methnani J, Kanzali MM, Koukane A, Bedoui MH, Latiri I. Prevalence and risk factors of functional gastrointestinal disorders in early period medical students: a pilot study in Tunisia. *Libyan Journal of Medicine.* 2022, Oct 14;17(1).
25. Oshima T. Functional dyspepsia: Current understanding and future perspective. *Digestion.* 2024, Jan 8;105(1):26-33.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/119032>