

Factors Contributing to Obesity among Adolescents

Sunita Poudel^{a*} and Jaya Laxmi Pradhan^b

^a School of Health and Allied Sciences, Pokhara University, Pokhara, Nepal.

^b The Central Department of Home Science, T.U, Bagbazar, Kathmandu, Nepal.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100054>

Received: 04/04/2023

Accepted: 08/06/2023

Published: 29/06/2023

Original Research Article

ABSTRACT

Background: Obesity is a medical condition characterized by an abnormal fat accumulation which is detrimental to health. The role of infectious disease is diminishing and that of non-communicable disease are increasing rapidly in developing countries like Nepal. Adolescents are vulnerable in terms of nutrition and the obese adolescent turns into an obese adult which leads to reduction of productivity of nation.

Objectives: To determine the prevalence and knowledge of overweight/obesity among adolescents and evaluate the associated factors in Bharatpur Metropolitan Municipality of Chitwan district.

Methodology: A descriptive cross-sectional study was conducted where two secondary schools of Bharatpur Metropolitan Municipality were randomly selected by lottery method. All the students of grade 9 and 10 of those two schools were included in the study (Sample size: 304). The question was self-administered and; Body Mass index (BMI) status and Central obesity was calculated. Data was analyzed by using IBM SPSS version 26.

Major Findings: In total, 187 (61.5%) of the respondents had knowledge about obesity, 251 (82.6%) had junk foods eating habits, 28 (9.2%) did not involve in any physical exercise and 50.3% of the respondents spent their leisure time playing/watching gadgets. The prevalence rate of overweight and obesity was 10.2% and 4.9% respectively. The study confirms 3 (1.9%) of boys and 12 (8.1%) of girls were central obese. Further, the mean BMI score was 18.57 kg/m² (Mean ± SD: 18.57±2.56) and mean waist circumference for boys and girls was 72.25 cm and 69.59 cm respectively. The study shows significant association (p value<0.05) between father's occupation and mother's education with types of snacks provided. Furthermore, respondent's sex, meal

*Corresponding author: E-mail: poudelsunita28@gmail.com;

frequency and gadgets at home show statistically significant association (p value <0.05) with nutritional status of respondents.

Conclusion: The study suggests that preventive approaches should be followed from the very beginning to avoid the obesity and health problem. The food preferences of respondents, foods they consumed, lack of physical exercise, preference of junk foods, parental history of obesity, use of gadgets and online games would lead to serious health effects in the future.

Keywords: Obesity; overweight; adolescents; nutritional habits; BMI.

1. INTRODUCTION

Obesity is a medical condition characterized by an abnormal fat accumulation which is detrimental to health. According to World Health organization, obesity is defined as being at or above the 95th percentile of body mass index for age and sex and Overweight as being between the 85th and 95th percentiles of body mass index for age and sex. WHO defines Body mass index (BMI) as being weight in kilogram per height in square meter recommended for use in children and adolescent [1] Obesity is a common, serious and costly disease of adults, adolescents and children [2].

Worldwide obesity has nearly tripled since 1975. The prevalence of overweight and obesity among children and adolescents aged 5-19 has risen dramatically from 4% in 1975 to over 18% in 2016. The rise has occurred similarly among both boys and girls: in 2016 18% of girls and 19% of boys were overweight [1]. More than 1 billion people worldwide are obese and among them over 340 million are adolescents [3]. The issue has grown to epidemic proportions, with over 4 million people dying each year as a result of being overweight or obese in 2017. Most of the world's population lives in countries where overweight and obesity kills more people than underweight [1].

The proportion of women who were overweight or obese increased from 9% in 2006 to 13% in 2011 and 22% in 2016 [4]. However, the prevalence of overweight among children is steady at 1% since 1996 to 2022 [5]. Obesity and central obesity are associated with an increased risk of multiple chronic diseases, including diabetes, cardiovascular disease (CVD), hypercholesterolemia, asthma and cancer [6,7]. Previous studies using Nepal Demography and Health Survey (NDHS) 2016 showed variation in underweight and overweight and/or obesity by individual level factors, i.e., women compared to men, urban residents compared to rural residents, and wealthy individuals were more likely to be overweight and/or obese [8].

Although individual characteristics play an important role in manifesting health outcomes, including weight status, recent evidence suggest that health is also determined by population level characteristics such as residence, neighborhood's walk ability, availability of food and so on [9,10].

The adolescents' population in Nepal covers 24% of total population [11] and young people (age group 16-25 years) accounts for approximately 20.8% of total population [12]. Adolescence is a critical phase for the development of obesity because of various biological, psychological, social and environmental change [13] and adolescence overweight and obesity may persist into adulthood [14,15]. Overweight and obese adolescents also suffer short-term health consequences and may increase the risk of developing NCDs at a younger age and consequently a premature death [16,17]. In addition; their participation in school and other daily activities is also limited depending on the status of obesity [15,18].

2. OBJECTIVES OF THE STUDY

2.1 General Objective

The general objective of the study is to study the knowledge about health problems of obesity, determine the prevalence of obesity rate, factors related with obesity and nutritional status among school going adolescents of Bharatpur Metropolitan Municipality of Chitwan District.

2.2 Specific Objectives

- i. To find out the level of knowledge about health problems regarding obesity among adolescents
- ii. To find out the prevalence rate of overweight and obesity among school going adolescents
- iii. To explore the factors related with obesity among school going adolescents
- iv. To assess the association between factors associated with obesity in terms of nutritional status.

3. MATERIALS AND METHODS

3.1 Study Design, Sample Size and Population

A descriptive cross-sectional study was conducted among adolescents of grade 9 and 10 of two schools of Bharatpur Metropolitan Municipality of Chitwan district which is the district headquarter and fourth largest city of Nepal. Bharatpur is a major destination for higher education, health care and transportation in the region. The two schools (one government and one private) were selected by lottery method among all schools of Bharatpur and census method was done to collect information among those selected schools. The total sample was 304.

3.2 Sampling Methods

Bharatpur Metropolitan Municipality was selected purposively. There were a total of 226 schools in Bharatpur metropolitan municipality and among them 125 were governmental and 101 were private schools. Further, there were 60 primary schools, 46 lower secondary schools, 81 secondary schools and 39 higher secondary schools. Then, the 81 secondary schools were listed and among them, two schools were randomly chosen through a lottery method. All the students of grade 9 and 10 of those two schools were included in the study.

3.3 Data Collection Tools and Techniques

Information was collected using self-administered questionnaire. Questionnaire consists of different sections and they are: socio-demographic, health related, dietary habits, behavioral, lifestyle and nutritional status. Height, weight and waist circumference of the respondents were measured by the researcher.

For measuring height, respondent was made to stand bare feet on a floor against a wall and with feet parallel and joined together and with heels and buttock touching the wall. It was made sure that head was erect. The height was marked on the wall with the help of a ruler and marker and it was measured with the measuring tape. For measuring the weight, bathroom weighing machine was used. Before using, it was made sure that it is at zero error.

For measuring waist circumference, the respondents were made to stand straight. The midpoint of the top of the hip bone and bottom of the rib cage was found. The measuring tape was

used and the waist circumference was measured horizontally. The tape should be snug and not very tight. The Body Mass Index and Central obesity was calculated.

3.4 Statistical Analysis

Data entry and analysis was done through IBM SPSS Statistics Version 26. Obesity was measured by Body Mass Index (BMI) which is defined as the weight in kilograms divided by the square of the height in meter (kg/m^2). BMI level was calculated by percentile ranking for adolescents by plotting the BMI of each adolescent on BMI-for-age growth chart of World Health Organization (for either girls or boys). The value is mentioned below:

Underweight: Less than 5th percentile

Normal / Healthy weight: 5th percentile to less than 85th percentile

Overweight: 85th to less than 95th percentile

Obese: Equal to or greater than 95th percentile

Central obesity was calculated depending upon the waist circumference i.e. circumference ≥ 94 cm in male and ≥ 80 cm in female.

The age of the participants was calculated using date of birth reported by adolescents and date of data collection. Descriptive and inferential analysis was done through SPSS. The p-value below 0.05 was considered as statistically significant.

4. RESULTS AND DISCUSSION

4.1 Socio-Demographic Characteristics of the Respondents

The socio demographic characteristics of the respondents were shown in Table 1. Among a total of 304 students, the mean age of the respondents was 15 years. The mean age of adolescents in this study was similar to the study conducted on Kaski which shows (15.08) years [19]. The study shows that 156 (51.3%) were boys and 148 (48.7%) girls participated in the study. Further, the study shows that 162 (53.3%) were from Government schools and 142 (46.7%) were from private schools. In a study conducted in 2022 shows 75.1% of overweight respondents belongs to private schools [20].

Furthermore, 225 (74%) students lived in nuclear family while only 79 (26%) lived in joint family which is similar to the study conducted in Gokarneshwor municipality i.e 75.63% nuclear and 24.37% joint family [9].

Table 1. Socio-demographic characteristics of the respondents

		n=304	
Variables		Frequency	Percentage (%)
Age	Early adolescents	6	2
	Mid adolescents	184	60.5
	Late adolescents	114	37.5
	Mean age: 15 years		
Sex	Boys	156	51.3
	Girls	148	48.7
Family type	Nuclear	225	74
	Joint	79	26
Family income/month	≤20000	11	3.6
	>20000	246	80.9
	Don't know	47	15.5
Father's education status	Illiterate	23	7.6
	Formal education	19	6.3
	Primary	64	21.1
	Secondary	80	26.3
	Bachelors and above	106	34.9
Mother's education status	Illiterate	10	3.3
	Formal education	79	26
	Primary	59	19.4
	Secondary	77	25.3
	Bachelors and above	66	21.7
	Don't know	13	4.3
Mother's occupation	Agriculture	16	5.3
	Business	42	13.8
	Services	7	2.3
	Labour	1	0.3
	Foreign employment	3	1
	Housewife	235	77.3
Father's occupation	Agriculture	30	9.9
	Business	61	20.1
	Services	39	12.8
	Labour	47	15.5
	Foreign employment	49	16.1
	Others	78	25.7
Type of school	Government	162	53.3
	Private	142	46.7

4.2 Food Consumption Pattern of the Respondents

Table 2 of the study shows the food consumption pattern of the respondents. It shows that all of the respondents (304) consume cereals every day, 7 (2.3%) were vegetarians, 212 (69.7%) consume soft drinks daily and 102 (33.6%) consume junk foods every day. Karki et. al., mentioned 83.1% of the respondents use soft drinks [21] and Chaulagain shows 10.93% of the respondents as vegetarian [9].

4.3 Exercise Behavior of the Respondents

Table 3 represents the exercise behavior of the respondents which shows 276 (90.8%) of the respondents did physical exercise and among them 182 (59.9%) of the respondents performed physical exercise for less than 1 hour, 96 (31.6%) for 1-2 hours and 1 (0.3%) for 2-3 hours a day. The study conducted by Jain and et al. shows 12.4% of the respondents were not involved in any moderate physical activity [22]

which is higher compared to this study with 31.12 % of the respondents who performed moderate physical activity for greater than 1 hour which is similar to this study.

Table 2. Food consumption pattern of the respondents

n=304

Food Groups	Frequency of intake			
	Everyday	Twice a week	Sometimes	Never
Cereals	304(100%)	-	-	-
Roots and tubers	166(54.6)	70(23%)	41(13.5%)	9(3%)
VegeTables	303(99.7%)	-	-	1(0.3%)
Fruits	154(50.7%)	62(20.4%)	88(28.9%)	-
Meat	24(7.9%)	125(41.1%)	138(45.4%)	17(5.6%)
Eggs	35(11.5%)	98(32.2%)	164(53.9%)	7(2.3%)
Fish/sea food	6(2%)	71(23.4%)	167(54.9%)	60(19.7%)
Legumes, nuts and seeds	78(25.7%)	123(40.5%)	103(33.9%)	-
Milk and milk products	153(50.3%)	84(27.6%)	50(16.4%)	3(1%)
Oils and fats	304(100%)	-	-	-
Condiments	212(69.7%)	61(20.1%)	30(9.9%)	1(0.3%)
Soft drinks	21(69.7%)	61(20.1%)	30(9.9%)	1(0.3%)
Junk foods	102(33.6%)	37(12.2%)	112(36.8%)	53(17.4%)

Table 3. Exercise behavior of the respondents

n=304

Variables		Frequency (N)	(%)
Physical exercise	Yes	276	90.8
	No	28	9.2
Types of physical exercise *	Walk	108	35.5
	Run	84	27.6
	Cycling	125	41.1
	Yoga	56	18.4
	Indoor physical exercise	69	22.7
	Taekwondo	1	0.3
Hours spent/day	<1 hour	182	59.9
	1-2 hour	96	31.6
	2-3 hour	1	0.3

*Multiple response




Fig. 1. Knowledge on health problems of obesity

4.4 Knowledge on Health Problems of Obesity

The knowledge on health problems related to obesity of was shown in Fig. 1 which shows 62% of them had knowledge whereas 38% did not have any knowledge about health problems of obesity. Majority of the respondents (21.1%) mentioned diabetes as the health problem of obesity. A study in Southern Ethopia shows adolescents with poor nutritional knowledge were significantly associated with overweight/obesity [23].

4.5 BMI Status of the Respondents

The Fig. 2 shows BMI status of respondents with prevalence of overweight of 10.2% and obesity (4.9%). The mean BMI was 18.57 kg/m². It was measured according to the WHO BMI calculation for age 5 to 19 years. The findings of the study conducted in Tulsipur municipality shows that prevalence of overweight and obesity was 4.25% and 2.25% which is low compared to this study [6] but is similar to the study conducted by Piryani et.al. i.e. 12.2% [15].

Further, a study conducted in Kaski, Nepal shows similar findings i.e 10% overweight and 3.3% obese [24]. Further, the prevalence was less in the study conducted by Sah et al. in Biratnagar i.e Overweight (2.9%) and Obese (1.8%) [25]. In an observational study conducted in adolescents between the ages 11 to 19 years in Bangalore, India , the prevalence of obesity was 5.9% [13]. The findings on the Prevalence of Obesity of this study is more than the findings of study on risk factors of non-communicable diseases conducted in Ratnanagar Municipality of Chitwan District which is 5.8% [22] and the findings of study conducted in Islamabad, Pakistan i.e. 8.5% [26]. Also, the findings of prevalence of overweight was lesser to some extent then study conducted by STEP survey of Nepal which is 17.7% [27], study conducted in rural china which is 24.8% [28], study conducted in Shenyang china which is 14.6% [29] and study conducted among 7-17 years old children and adolescents in Lithuania is 12.6% [30]. The dissimilarity in findings may be because of variation in level of socio-economic development and lifestyle differences in various part of world.




Fig. 2. BMI status of respondents

4.6 Central Obesity among Respondents

Fig. 3 represents the central obesity of respondents which shows that 3 (1.9%) of boys were central obese and 12 (8.1%) of girls were central obese. It shows that girls were more than four times central obese than boys. While measuring the waist circumference of the respondent, they are central obese if waist circumference is greater or equals to 94 cm in boys and greater or equals to 80 cm in female. In the study conducted by Castro et al. in Brazil shows the prevalence of abdominal obesity was 10.6% for the total sample (10.5% male, 10.8% female) [31]. The reasons for higher obesity among girls may be because of their fewer involvements in sports and other games. And difference in findings among above study may be because of difference in study site.




Fig. 3. Central obesity of respondents

4.7 Association between parents' education and occupation with types of food provided

Table 4 shows that the food items provided to the adolescents is statistically significant with the educational status of mother. The mothers usually serve food items to the children and their educational status directly depends upon the types of food items they provided to their children. Most of the educated mothers provide healthy foods to their children. Further, it shows the significant association between occupation of father and food items the respondents' are provided as snacks. It shows most of the children whose father had a business consumed junk foods as snacks. This study contrasts to the study conducted in urban Nepal which shows the greater likelihood of being overweight or obese of children whose mothers in a professional field (aOR = 1.34, 95% CI: 1.02–4.05) [21] and study conducted in Dang which shows significant association between BMI and occupation of mothers [32]. In addition, the study conducted in Lithuania showed statistical significant with obesity and education of parents [30] and the study conducted in 20 Italian region also showed significant association with obesity [14].

4.8 Association between Gadgets Use and Nutritional Status

The findings of this study (Table 5) shows significant association (p value: 0.03) with gadgets at home of the respondents and nutritional status however no association was found with duration of gazettes use by the respondents. This findings contrast with the findings of the study conducted among urban school students of Lalitpur sub-metropolitan city that showed watching television for more than 2 hrs per day had association with obesity [15]. In addition, the study conducted by Regmi and Purushothama in Tulsipur municipality, Dang shows that higher screening time for TV/Computer had association with obesity [6].

4.9 Association between obesity and type of school

The Table 6 shows significant association between obesity and type of school (p value=0.03) which shows students studying in private schools are more obese compared to the government schools. A study by Sah et al. show students studying in private schools and family income > Rs.10,000 were strongly associated

with overweight and obesity [25] and the study shows significant association i.e. $p < 0.05$ conducted in Tulsipur municipality, Dang also between obesity and school type [6]

Table 4. Association between parents' education and occupation with food types

n=304

Education/ Occupation status		Food items (Snacks)			χ^2	P value
		Home made	Healthy canteen foods	Junk foods		
Education-Father	Illiterate	1	2	20	8.64	0.56
	Formal education	0	1	10		
	Primary	3	0	34		
	Secondary	9	5	65		
	Bachelors and above	11	10	70		
	Don't know	2	1	8		
Education-Mother	Illiterate	1	3	6	23.9	0.008*
	Formal education	3	1	53		
	Primary	2	1	37		
	Secondary	12	5	59		
	Bachelors and above	5	8	44		
	Don't know	3	1	8		
Occupation-Father	Agriculture	2	2	26	26.9	0.008*
	Business	3	6	44		
	Services	2	5	17		
	Labour	1	0	27		
	Foreign employment	8	1	40		
	Others	7	1	43		
	Don't know	3	4	10		
	Occupation-Mother	Agriculture	1	3		
Business	1	2	31			
Services	0	2	4			
Labour	0	0	1			
Foreign employment	0	0	3			
Others	20	9	142			
Don't know	4	3	14			

* Significant association i.e. $p \text{ value} < 0.05$

Table 5. Association between gadgets use and nutritional status

n=304

Variables		Nutritional status				χ^2	P value
		Under weight	Normal weight	Overweight	Obese		
Gadgets at home	Yes	15	228	25	14	8.37	0.03*
	No	0	15	6	1		
Time spent on gadgets	less than 1 hour	4	92	7	6	7.89	0.54
	1-2 hours	6	102	13	8		
	2-3 hours	2	13	2	0		
	>3 hours	3	21	3	0		

* Significant association i.e. $p \text{ value} < 0.05$

Table 6. Association between obesity and type of school

n=304

School type	Obesity		χ^2	P value
	Present	Absent		
Government	3	159	7	0.03*
Private	11	131		

* Significant association i.e. p value < 0.05

5. CONCLUSION

In conclusion, the study shows that there is high prevalence of overweight among adolescents and they are at more risk of getting obese. The occupation of parents and educational status is directly associated with the types of food provided. The gadgets at home and school types show positive association with nutritional status in the study. The food preference of the respondents, the food groups they consumed, lack of physical exercise, preference of junk foods and juices, parental history of obesity, addiction to the gadgets and online games would lead to serious health effects in the future as well as in their academic performances.

The recommended measures from the study are:

- Timely identification and prevention of overweight/obesity is required for prevention of development of other co-morbidities.
- Healthy foods should be provided in home and school canteen strictly prohibiting junk foods.
- Students should be encouraged to do physical activities like walk, yoga, exercise, outdoor games etc.
- Program regarding weight management should be conducted for both teachers and students.
- Multi-sectoral approach for the prevention and control of overweight and obesity.
- An enabling environment is of paramount importance to increase awareness about the risk factors for obesity in adolescence to decrease the prevalence of obesity associated NCDs in the upcoming generations of Nepal.

CONSENT AND ETHICAL APPROVAL

The ethical approval was obtained from Central Department of Home Science, Bagbazar, Kathmandu. The study was conducted with prior permission from the school authorities. Before data collection, the research purpose and process of the study was clearly explained.

Verbal informed consent was taken from participants before administering questionnaires. The study participants were assured for confidentiality of the information at all levels of the study by avoiding personal identifiers. The respondent may leave whenever they want. After the data collection, the researcher explained the school adolescents about overweight and obesity, their causes, consequences, and preventive measures.

ACKNOWLEDGEMENTS

We are grateful to the Central Department of Home Science, Kathmandu for the opportunity and so thankful to Mr. Narayan Gautam and Mr. Navaraj Poudel for their tremendous support, help and cooperation throughout the data collection period. We are thankful to all the study participants who provided their valuable time and cooperation in this venture.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Obesity and overweight [Internet]. [cited 2023]. Available: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
2. CDC Overweight & Obesity [Internet]. Centers for Disease Control and Prevention. 2023 [cited 2023]. Available: <https://www.cdc.gov/obesity/index.html>
3. World Obesity Day 2022 – Accelerating action to stop obesity [Internet]. [cited 2023]. Available: <https://www.who.int/news/item/04-03-2022-world-obesity-day-2022-accelerating-action-to-stop-obesity>
4. MOH/Nepal M of H, ERA/Nepal N, ICF. Nepal Demographic and Health Survey 2016. 2017 Nov 1 [cited 2023];

- Available:<https://dhsprogram.com/publications/publication-fr336-dhs-final-reports.cfm>
5. Nepal Health Facility Survey Final Report. 2021.
 6. Regmi S, Purushothama J. Overweight and obesity among higher secondary level students of Tulsipur Municipality, Nepal. *Int J Community Med Public Health*. 2018; 24;5:774.
 7. Nonboonyawat T, Pusanasuwannasri W, Chanrat N, Wongthanavimok N, Tubngern D, Panutrakul P, et al. Prevalence and associates of obesity and overweight among school-age children in a rural community of Thailand. *Korean J Pediatr*. 2019;62(5):179–86.
 8. Prevalence and factors affecting underweight, overweight and obesity using Asian and World Health Organization cutoffs among adults in Nepal: Analysis of the Demographic and Health Survey 2016 - ScienceDirect [Internet]. [cited 2023]. Available:<https://www.sciencedirect.com/science/article/abs/pii/S1871403X18303533>
 9. Chaulagain K. The Burden of Overweight and its Risk Factors in Urban Nepalese School Adolescents: A Cross- Sectional Study. *J Qual Health Care Econ*. 2021; 12;4.
 10. Shrestha N, Mishra SR, Ghimire S, Gyawali B, Pradhan PMS, Schwarz D. Application of single-level and multi-level modeling approach to examine geographic and socioeconomic variation in underweight, overweight and obesity in Nepal: findings from NDHS 2016. *Sci Rep*. 2020;12;10(1):2406.
 11. World Population Day [Internet]. UNFPA Nepal. 2016 [cited 2023]. Available:<https://nepal.unfpa.org/en/node/10804>
 12. For young people, by young people [Internet]. UNFPA Nepal. 2022 [cited 2023]. Available:<https://nepal.unfpa.org/en/news/young-people-young-people>
 13. Kamath SV Sangamesh G. Prevalence of obesity and its risk factors in school going adolescents of urban Bangalore, India. *Int J Contemp Pediatr*. 2016;28;3(2):568–74.
 14. Lazzeri G, Giacchi MV, Spinelli A, Pammolli A, Dalmaso P, Nardone P et al. Overweight among students aged 11–15 years and its relationship with breakfast, area of residence and parents' education: results from the Italian HBSC 2010 cross-sectional study. *Nutr J*. 2014;5;13:69.
 15. Piryani S, Baral KP, Pradhan B, Poudyal AK, Piryani RM. Overweight and its associated risk factors among urban school adolescents in Nepal: a cross-sectional study. *BMJ Open*. 2016;20;6(5): e010335.
 16. Ranjani H, Pradeepa R, Mehreen TS, Anjana RM, Anand K, Garg R, et al. Determinants, consequences and prevention of childhood overweight and obesity: An Indian context. *Indian J Endocrinol Metab*. 2014;18(Suppl 1): S17–25.
 17. Raj M, Kumar RK. Obesity in children & adolescents. *Indian J Med Res*. 2010; 132(5):598–607.
 18. Story M, Nannery MS, Schwartz MB. Schools and Obesity Prevention: Creating School Environments and Policies to Promote Healthy Eating and Physical Activity. *Milbank Q*. 2009;87(1):71–100.
 19. (PDF) Prevalence and Knowledge on Obesity Among School Going Adolescents of Kaski, Nepal [Internet]. [cited 2023]. Available:https://www.researchgate.net/publication/330334109_Prevalence_and_Knowledge_on_Obesity_Among_School_Going_Adolescents_of_Kaski_Nepal
 20. Singh DR, Sunuwar DR, Dahal B, Sah RK. The association of sleep problem, dietary habits and physical activity with weight status of adolescents in Nepal. *BMC Public Health*. 2021;17;21(1):938.
 21. Deepak Karki. The Basics of Monitoring, Evaluation and Supervision of Health Servic... [Internet]. 22:57:19 UTC [cited 2023 Jan 24]. Available:<https://www.slideshare.net/dekarki/the-basics-of-monitoring-evaluation-and-supervision-of-health-services-in-nepal-37391386>
 22. Adhikari K, Jain V, Adak M, Gupta N, Koshy A. Prevalence of Risk Factors of Non-Communicable Diseases among Adolescent in Parsa District of Nepal. 2013.
 23. Kedir S, Hassen K, Melaku Y, Jemal M. Determinants of overweight and/or obesity among school adolescents in Butajira Town, Southern Ethiopia. A case-control study. *PLOS ONE*. 2022;28;17(6): e0270628.
 24. Pandey A, Sapkota S. Prevalence and Knowledge on Obesity among School Going Adolescents of Kaski, Nepal. *J Nepal Paediatr Soc*. 2019;11;38: 63–8.

25. Sah VK, Giri A, Acharya R. Prevalence of overweight, obesity and its associated risk factors among school children aged 6-16 years of Biratnagar. *J Nobel Med Coll.* 2016;26;5(2):22–5.
26. Department of Public Health and Community Medicine, AJK Medical College, University of Azad Jammu and Kashmir, Muzaffarabad, Azad Kashmir, Durrani HM, Durrani SM, Alumni-Rawalpindi Medical College, Tipu Rd., Rawalpindi, Pakistan, Kumar R, Health Services Academy, Islamabad, Pakistan, et al. Prevalence of Overweight and Obesity Among Adolescents of Islamabad: A Cross Sectional Study. *Pediatr Neonatal Nurs - Open J.* 2017;3;4(1):8–11.
27. Findings of prevalence of overweight by STEP survey of Nepal (Step Survey, 2013 - Google Search [Internet]. [Cited 2023].
28. He Y, Pan A, Yang Y, Wang Y, Xu J, Zhang Y, et al. Prevalence of Underweight, Overweight, and Obesity Among Reproductive-Age Women and Adolescent Girls in Rural China. *Am J Public Health.* 2016;106(12):2103–10.
29. Zhai L, Dong Y, Bai Y, Wei W, Jia L. Trends in obesity, overweight, and malnutrition among children and adolescents in Shenyang, China in 2010 and 2014: a multiple cross-sectional study. *BMC Public Health.* 2017;2;17(1):151.
30. Smetanina N, Albaviciute E, Babinska V, Karinauskiene L, Albertsson-Wikland K, Petrauskiene A, et al. Prevalence of overweight/obesity in relation to dietary habits and lifestyle among 7-17 years old children and adolescents in Lithuania. *BMC Public Health.* 2015;1;15:1001.
31. Castro JAC, Nunes HEG, Silva DAS. Prevalence of abdominal obesity in adolescents: association between sociodemographic factors and lifestyle. *Rev Paul Pediatr.* 2016;34(3):343–51.
32. Bhattarai S, Bhusal CK. Prevalence and associated factors of malnutrition among school going adolescents of Dang district, Nepal. *AIMS Public Health.* 2019;21;6(3):291–306.

© 2023 Poudel and Pradhan; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/100054>