



Effect of Plant Growth Regulators at Different Levels of Zinc on Growth and Yield of Groundnut (*Arachis hypogea* L.)

Kakulavarapu Navya Sri Subha^{a*} and Biswarup Mehera^a

^a Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj - 211007, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i1030831

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/87096>

Original Research Article

Received 05 March 2022

Accepted 10 May 2022

Published 19 May 2022

ABSTRACT

The field experiment entitled "Effect of Plant Growth Regulator at different levels of Zinc on growth and yield of Groundnut (*Arachis hypogea* L.)" was conducted during *kharif* season of 2021 at Crop Research Farm, Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P). The experiment plot was sandy loam in texture, nearly acidic in soil reaction (PH 6.9), low in organic carbon (0.112%), available N (278.93Kg/ha), available P(10.8Kg/ha) and available K (206.4Kg/ha).The experiment was laid out in Randomized Block Design, treatments of different plant growth regulator levels (75, 100 and 125 ppm) and zinc levels viz.(20,25 and 30 kg/ha) were imparted in the experiment and were replicated thrice.

The experiment was laid out in Randomized Block Design comprised of 3 replications and total 9 treatments viz. T1: (GA3)75PPM + 20 kg/ha ZnSO₄, T2: (GA3) 75PPM + 25 kg/ha ZnSO₄, T3: (GA3) 75PPM+ 30 kg/ha ZnSO₄, T4:(GA3) 100PPM + 20 kg/ha ZnSO₄, T5: (GA3) 100PPM +25 kg/ha ZnSO₄, T6: (GA3) 100PPM + 30 kg/haZnSO₄, T7: (GA3) 125PPM + 20 kg/ha ZnSO₄, T8: (GA3) 125PPM + 25 kg/ha ZnSO₄, T9: (GA3) 125PPM +30 kg/ha ZnSO₄. Results obtained indicated that there was significant increase in growth parameters at harvest viz., plant height (58.40cm), number of nodules/plant (59.77), dry weight accumulation(38.1g),and yield attributes viz., No.pods/plant(19.3), No.kernals/pod (4.3), seed index(44.50), seed yield (3169 kg/ha), Haulm

yield (4747.3kg/ha) were recorded with the application of (GA3125ppm+ ZnSO₄ 30kg/ha). However, Maximum Gross return (142636.5 INR/ha), Net return (102971.1INR/ha) and B:C ratio (2.59) was recorded with the application of (GA3125ppm+ ZnSO₄ 30kg/ha). Therefore, it is concluded that the application of (GA3125ppm+ZnSO₄30kg/ha) was more productive and economically feasible.

Keywords: Groundnut; plant growth regulator (GA3); zinc; economics.

1. INTRODUCTION

Groundnut (*Arachis hypogea* L.) is one of the commercial crops of the world that rank 13th among the food crops grown over an area of 5.40 million ha and production of 5.43 million tonnes with a productivity of 910kg/ha (AICRP 2020). About two third of the world's groundnut production is used as oil and remaining one third is consumed as food. Being the best source of vegetable oil, it also provides raw material for industrial serving as concentrated animal feed on organic manure. It contributed to sustainable agriculture being a legume and cultivated since long in both kharif and summer season by the agriculturist. Groundnut contains on an average 12-15 percent carbohydrate, 25-30 percent protein and 45-50 percent oil. The high-energy value protein content and minerals make groundnut a rich source of nutrition at a comparatively low price.

Among plant growth regulators, Gibberellin play a vital role in regulating developmental processes within plant bodies. Gibberellin helps in cell growth of stem, leaves and other aerial parts b causing cell elongation, and increase in intermodal length. A higher concentration of gibberellins increases plant growth [1]. The different concentrations of GA had significant effect on growth in mustard [2]. GA3 is an important growth regulator that may have many uses to modify the growth, yield and yield attributing characters of plant. Plant growth regulators are used widely to improve plant performance. GA3 is one of those growth regulators that have positive effect on plant as enhancing vegetative growth and plant yield and increasing dry weight [3]. GA3 has been widely used to improve germination rate, seedling growth and consequently yield. Yakubu et al., [4], Rastogi et al., [5].

In India, zinc is now considered as the fourth most important yield limiting nutrient after nitrogen, phosphorus and potassium, respectively. Among oilseeds, groundnut in

particular suffers from Zn deficiency [6]. Zinc deficiency start yellowing of leaves from lamina to base, mid-rib and veins remain green, later on necrotic brown spots are developed and dorsal leaf veins become brown. It is likely to occur when soils are low in organic matter and alkaline under high levels of soil p and soils are cool and wet during the vegetative phase, zinc increases the chlorophyll content in the leaves and nodule number, pod production and weight. Zinc's function in cell membrane integrity will also be discussed especially for root cells along with its role in suppressing free radical damage to cells [7]. Zinc increases the resistance of plants to pathogens by bringing changes in anatomy and physiology of host plant.

2. MATERIALS AND METHODS

The experiment was carried out during *Kharif* season of 2021 at the CRF (Crop Research Farm) SHIATS, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, Uttar Pradesh. The crop Research Farm is situated at 25.75⁰ N latitude, 87.19⁰ E longitude and at an altitude of 98m above mean sea level. This area is situated on the right side of the river Yamuna and by the opposite side of Prayagraj City. All the facilities required for crop cultivation were available. The experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 6.9), low in organic carbon (0.112%), available N (278.93 Kg/ha), available P (10.8 Kg/ha) and available K (206.4 Kg/ha). The crop was sown on 20th July 2020 using variety Kadiri-6. The experiment was laid out in Randomized Block Design comprised of 3 replications and total 9 treatments viz. T1: (GA3) 75PPM + 20 kg/ha ZnSO₄, T2: (GA3) 75PPM + 25 kg/ha ZnSO₄, T3: (GA3) 75PPM+ 30 kg/ha ZnSO₄, T4: (GA3) 100PPM + 20 kg/ha ZnSO₄, T5: (GA3) 100PPM +25 kg/ha ZnSO₄, T6: (GA3) 100PPM + 30 kg/ha ZnSO₄, T7: (GA3) 125PPM + 20 kg/ha ZnSO₄, T8: (GA3) 125PPM + 25 kg/ha ZnSO₄, T9:

(GA3) 125PPM +30 kg/ha ZnSO₄. All nutrients were applied into the soil in the form of Urea, Single super phosphate (SSP) and Muriate of potash (MOP). Entire dose of P and K was applied basal for respective plots, half dose of N (as urea) was applied as basal, one-fourth at 30 days after sowing and remaining one-fourth at the time of flowering. Zinc levels are (20,25,30 kg/ha) was applied as soil application along with NPK fertilizers before sowing. Plant growth regulator levels are (75,100,125 ppm) was applied as foliar spray through Knapsack sprayer as two doses. One at 22 DAS and another one at 45DAS. The growth parameters were recorded at periodical intervals of 15,30,45,60,75 DAS and at harvest stage from the randomly selected five plants in each treatment. Statistically analysis was done for all the parameters in one way Anova and mean compared at 5% probability level of significant results.

3. RESULTS AND DISCUSSION

3.1 Effect of Plant Growth Regulator at Different Levels of Zinc on Growth Parameters of Groundnut

Effect of Plant Growth Regulator and Zinc on growth parameters of Groundnut of are presented in Table 1. In the results revealed that maximum plant height (58.40cm), number of nodules/plant(59.77), dry weight(38.1), was observed with application of (GA3) 125PPM + 30Kg/ha ZnSO₄ where crop growth rate(10.00) and relative growth rate(0.009) was observed with application of(GA3) 125PPM + 30Kg/ha ZnSO₄ as closely followed by the Plant height (56.86), number of nodules/plant(58.40), Dry matter accumulation (37.3) which was recorded in the treatment of ((GA3) 125PPM + 25Kg/ha ZnSO₄). Maximum plant height and maximum number of nodules are obtained due to the corresponding increase in level of GA3 at all the growth stages and the function of zinc is to produce chlorophyll to the plant, electron transportation, enzyme activation and Zinc can help in the nodulation activity due to enzymatic activity and nitrogen fixation which ultimately increase nodulation. Higher dry matter accumulation depends upon the photosynthesis and respiration rate, which finally increases the plant growth with respect to increased plant height, number of nodules etc. The favourable effect of plant growth regulator and zinc might influence the metabolism of the plant, effect on

photosynthetic pigments and activity of enzymes which in turn helps to increase in the vegetative growth. The results on higher crop growth was also reported by A Akter et al., [2] , Reddi Naveen kumar et al., [8].

3.2 Effect of Plant Growth Regulator at Different Levels of Zinc on Yield Parameters of groundnut

Effect of Plant Growth Regulator at different levels of Zinc on yield parameters of Groundnut are presented in Table 2. In the results revealed that higher No. pods/ plant (19.3), No. kernels/pod(4.3), Seed index (44.50), Seed yield (3169.7 kg/ha) and Haulm yield (4747.3 kg/ha) was recorded with application of (GA3) 125PPM + 30Kg/ha ZnSO₄ and Harvest Index (41.6) was observed with application of (GA3) 125PPM + 25Kg/ha ZnSO₄ as closely followed by the No. pods/ plant(18.8), No. kernels/pod(3.9), Seed index (41.40), Seed yield (2923.6kg/ha) was observed with the application of (GA3) 125PPM + 25Kg/ha ZnSO₄). The favourable combination of GA3 and zinc plays a vital role in increasing seed yield because they both takes place in many physiological processes like increasing the chlorophyll content, carbohydrate metabolism, protein synthesis, regulating the stomata, cell elongation, pod formation, increasing the flowering set which enhance the seed yield. The yield of a crop is the cumulative effect of yield attributing characters such as pods per plant, kernels per pod and seed index. Thus, the seed yield of ground nut also increased significantly due to application of zinc as a consequence of highest values of above parameters. The increase in seed yield due to zinc and GA3 application might be due to the cumulative effect of increased plant height, number of nodules per plant and dry matter production i.e., increased growth parameters. The seed and haulm yields combined together showed significant increase in biological yield of ground nut. The results of the present investigation are in close conformity with the findings of Sai Surya Goutami V. and Ananda N (2018).

3.3 Effect of Plant Growth Regulator at Different Levels of Zinc on Economics

Effect of Plant Growth Regulator at different levels of Zinc on economics of Groundnut

Table 1. Effect of Plant Growth Regulator at different levels of zinc on growth parameters at harvest of groundnut

Treatments	Plant height(cm)	No.of nodules per plant	Plant dry weight(g)	Crop growth rate(g/m ² /day)	Relative growth rate(g/g/day)
(GA ₃)75PPM+20Kg/ha Znso ₄	52.46	46.99	34.1	10.00	0.009
GA ₃)75PPM+25Kg/ha Znso ₄	53.50	48.77	34.8	7.41	0.006
GA ₃)75PPM+30Kg/ha Znso ₄	54.86	50.77	35.7	7.90	0.006
GA ₃)100PPM+20Kg/ha Znso ₄	54.53	49.55	35.1	7.53	0.006
GA ₃)100PPM+25Kg/ha Znso ₄	54.93	51.66	35.9	7.66	0.006
GA ₃)100PPM+30Kg/ha Znso ₄	55.86	53.99	36.6	7.29	0.005
GA ₃)125PPM+20Kg/ha Znso ₄	55.93	52.55	36.2	7.41	0.005
GA ₃)125PPM+25Kg/ha Znso ₄	56.86	58.40	37.3	6.91	0.005
GA ₃)125PPM+30Kg/ha Znso ₄	58.40	59.77	38.1	7.16	0.005
F test	S	S	S	NS	NS
SEm(±)	0.62	0.63	0.33	0.95	0.0009
CD(5%)	1.88	1.09	0.99	-	-

Table 2. Effect of Plant Growth Regulator at different levels of zinc on yield parameters of groundnut

Treatments	No.of plants/plant	No.of kernals per pod	Seed index(g)	Seed yield (Kg/ha)	Haulm yield(Kg/ha)	Harvest index(%)
(GA ₃)75PPM+20Kg/ha Znso ₄	15.5	2.7	39.40	2460.7	3788.7	38.6
GA ₃)75PPM+25Kg/ha Znso ₄	17.3	3.2	39.51	2536.9	3985.3	38.9
GA ₃)75PPM+30Kg/ha Znso ₄	17.8	3.6	40.23	2463.8	4145.3	37.5
GA ₃)100PPM+20Kg/ha Znso ₄	17.4	3.3	39.93	2547.3	3640.3	41.2
GA ₃)100PPM+25Kg/ha Znso ₄	18.3	3.6	40.37	2633.4	36749.0	41.3
GA ₃)100PPM+30Kg/ha Znso ₄	18.6	4.0	41.10	2676.6	4526.7	47.9
GA ₃)125PPM+20Kg/ha Znso ₄	18.5	3.8	40.93	2774.0	4378.0	38.8
GA ₃)125PPM+25Kg/ha Znso ₄	18.8	3.9	41.40	2923.6	4098.7	41.6
GA ₃)125PPM+30Kg/ha Znso ₄	19.3	4.3	44.50	3169.7	4747.3	40.0
F test	S	S	S	S	S	S
SEm(±)	0.22	0.14	1.08	13.06	130.36	0.73
CD(5%)	0.66	0.42	0.42	39.15	390.83	2.19

Table 3. Effect of Plant Growth Regulator at different levels of zinc on Economics of groundnut

Treatments	Cost of cultivation(INR/ha)	Gross returns(INR/ha)	Net return (INR/ha)	B:C ratio
(GA ₃)75PPM+20Kg/ha Znso ₄	38790.4	110731.5	71941.1	1.85
GA ₃)75PPM+25Kg/ha Znso ₄	38915.4	114160.5	75254.1	1.93
GA ₃)75PPM+30Kg/ha Znso ₄	39165.4	110871.0	71705.6	1.83
GA ₃)100PPM+20Kg/ha Znso ₄	38990.4	114628.5	75638.1	1.94
GA ₃)100PPM+25Kg/ha Znso ₄	39115.4	118503.0	79387.6	2.02
GA ₃)100PPM+30Kg/ha Znso ₄	39369.4	120312.0	80946.6	2.05
GA ₃)125PPM+20Kg/ha Znso ₄	39290.4	124830.0	85539.6	2.17
GA ₃)125PPM+25Kg/ha Znso ₄	39415.4	131562.0	92146.6	2.33
GA ₃)125PPM+30Kg/ha Znso ₄	39665.4	142636.5	102971.1	2.59

presented in Table 3. The highest gross returns (142636.51INR/ha), higher net returns (102971.11INR/ha) and maximum B:C ratio (2.59) recorded with the application of (GA3) 125PPM + 30Kg/ha ZnSO₄. This was mainly due to higher seed and haulm yields compared to other treatment combinations. Gross returns, net returns and B: C ratio increased significantly due to successive increase in varying levels of PGR and Zinc with application of (GA375ppm +ZnSO₄ 30Kg/ha). This might be attributed to higher seed and haulm yields obtained with comparatively less cost than additional income under these treatments. Similarly results were also reported by Chittam Ravi Kishore Reddy et al. [9].

4. CONCLUSION

It may be concluded that application of (GA3) 125PPM + 30Kg/ha ZnSO₄ showed higher yield and economics. Therefore, it is recommended for farmers for receiving higher yield and economic benefits of Groundnut [10-22].

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Bora RK, Sarma CM. Effect of Gibberellic acid and cycocel on growth, yield and protein content of pea. Asian Journal of Plant Sciences; 2006.
- Akter A, Islam E, Karim R, Razzaque AHM. Effect of GA3 on growth and yield of Mustard. International Journal of Sustainable Crop Production. 2007;2(2):16-20.
- Islam MR, Hasan M, Akter N, Akhtar S. Cytokinin and gibberellic acid alleviate the effect of waterlogging in mungbean. journal clean was. 2021;5:21-26.
- Yakubu H, Izge AU, Hussaini MA, Jibrin JM, Bello OG, Isyaku MS. Varietal response and gibberellic acid concentrations on yield and yield traits of groundnut (*Arachis hypogaea* L.) under wet and dry conditions. Academia journal of Agricultural Research. 2013;1 (1):001-008.
- Rastogi A, Siddiqui A, Mishra BK, Srivastava M, Pandey R, Misra P, Shukla S. Effect of auxin and gibberellic acid on growth and yield components of linseed (*Linum usitatissimum* L.). Crop Breeding and Applied Biotechnology. 2013;13:136-143.
- Singh AL. Prevention and correction of zinc deficiency of groundnut in India. In proceeding of zinc crops. Conference for improving crop production and human health, Istanbul, Turkey; 2007.
- Cakmak I. Tansely Review No.111. possible roles of zinc in protecting plant cells from damage by reactive oxygen speices. New phytologist. 2000;146(2):185-205.
- Reddi Naveen Kumar, Rajesh Singh, Wasim Khan. Effect of spacing and plant growth regulators on growth and yield of zaid sesame. International Journal of Current Microbiology and Applied Sciences. 2021;10(03):205-210.
- Reddy CRK, Dawson J, Talasila V. Biofortification of zinc with different phosphorous sources on growth and yield of groundnut (*Arachis hypogaea* L.). Journal of Pharmacognosy and Phytochemistry. 2019;8(3):4439-4442.
- Das S, Green A. Importance of zinc in crops and human health. J SAT Agric Res. 2013;11:1-7.
- Deo C, Khandelwal RB. Effect of zinc and phosphorus on yield, nutrient uptake and oil content of mustard grown on the gypsum-treated sodic soil. Journal of the Indian Society of Soil Science. 2009;57(1):66-70.
- Gowthami V, Ananda N. Impact of Zinc and Iron Ferti-Fortification on Leaf Area Index, Kernel Yield, Shelling Percentage and Iron Uptake of Groundnut (*Arachis hypogaea* L.) Genotypes. International Journal of Agriculture, Environment and Biotechnology. 2018;11(5):755-760.
- Irmak S, Cil AN, Yucel H, Zulkuf KAYA. Effects of Zinc Application on Yield and Some Yield Components in Peanut *Arachis hypogaea* in the Eastern Mediterranean Region. Journal of Agricultural Sciences. 2015;22(1):109-116.
- Tripathi MK, Chaturvedi S, Shukla DK, Saini SK. Influence of integrated nutrient management growth, yield and quality of Indian mustard (*Brassica juncea* L.) in tarai region of northern India. Journal of Crop Weed. 2011;7(2):104-107.
- Vekaria GB, Rakholiya KD, Vora VD, Patel JT, Sutaria GS, Vekaria PD. Response sesame (*Ses amum indicum* L.) to growth

- regulator under dry farming condition. Int J Curr Microbiol App Sci. 2017;6:1113-1120.
16. Verma CK, Prasad K, Yadav DD. Studies on response of sulphur, zinc and boron levels on yield, economics and nutrients uptake of mustard [brassicajuncea(l.) czern & coss.]. Crop Research. 2012;44(1and2):75-78.
 17. Woodroof JG. Peanuts: production, processing products; 1996.
 18. Reddi Naveen Kumar, Rajesh Singh, Wasim Khan. Effect of spacing and plant growth regulators on growth and yield of zaid sesame. International Journal of Current Microbiology and Applied Sciences. 2021;10(03):205-210.
 19. Singh AL. Prevention and correction of zinc deficiency of groundnut in India. In proceeding of zinc crops. Conference for improving crop production and human health, Istanbul, Turkey; 2007.
 20. Tripathi MK, Chaturvedi S, Shukla DK, Saini SK. Influence of integrated nutrient management on growth, yield and quality of Indian mustard (*Brassica juncea* L.) in tarai region of northern India. Journal of Crop Weed. 2011;7(2):104-107.
 21. Vekaria GB, Rakholiya KD, Vora VD, Patel JT, Sutaria GS, Vekaria PD. Response of sesame (*Ses amum indicum* L.) to growth regulator under dry farming condition. Int J Curr Microbiol App Sci. 2017;6:1113-1120.
 22. Verma CK, Prasad K, Yadav DD. Studies on response of sulphur, zinc and boron levels on yield, economics and nutrients uptake of mustard [brassicajuncea(l.) czern & coss.]. Crop Research. 2012;44(1and2):75-78.

© 2022 Subha and Mehera; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/87096>