



Search for Local Food Supplements: Study of the Nutritional Value of 15 Plant Species from Niger

Soumana Sourounda Ide ^{a*},
Mahamane Sabiou Sani Maazou ^a,
Saadatou Alkassoum Sangare ^b,
Maman Mahaman Salissou ^c,
Almoustapha Théodore Yatta ^d,
Mahamane Djika Hachimou ^e,
Garba dit Gado Halidou Ousseini ^a,
Almou Abdoulaye Alio ^a, Sabo Haoua Seini ^a
and Ameyapoh Yaovi ^f

^a Laboratory of Nutrition and Valorization of Agro-Resources, Department of Chemistry, Faculty of Science and Technology, Abdou Moumouni University, Niamey, Niger.

^b LARADES Laboratory-Djibo Hamani University of Tahoua/Public Policy Analysis and Government Action Evaluation Unit (CAPEG), Studies and Research Department, Niamey, Niger.

^c Ministry of Public Health, Population and Social Affairs, Niamey, Niger.

^d Ministry of Planning, National Institute of Statistics, National Information Platform for Nutrition Niamey, Niger.

^e Animal Production Laboratory, Faculty of Agronomy, Abdou Moumouni University, Niamey, Niger.

^f Laboratory of Microbiology and Quality Control of Foodstuffs, Higher School of Biological and Food Techniques, University of Lomé, Lomé, Togo.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJNFS/2024/v16i51427

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116415>

*Corresponding author: Email: soumanasouroundaide.97@gmail.com;

ABSTRACT

In Niger, the prevalence of malnutrition and mortality linked to undernutrition among Nigerien children under 5 years of age remains high and constitutes a major concern for public authorities. However, the country is teeming with significant biodiversity that could help combat high levels of undernutrition among children aged 6 to 59 months. The present study aimed to evaluate the nutritional quality of 15 plant species of which 17 samples were taken as potential complementary foods. The water, ash, lipids, protein and cellulose content were determined. Carbohydrate content and energy density were calculated. Biochemical characterization of the sampled foods showed that *Glycine max* seeds were richer in protein (36.39%). *Neocarya macrophylla* almond has a relatively high lipid content (60.75%) and cellulose content (26.74%). The pulp of *Hyphaene thebaica* was rich in mineral elements (7.16%). The tuber of *Ipomoea batatas* (95.25%) and the pulp of *Neocarya macrophylla* (93.01%) contained sufficient carbohydrates, and the humidity was found to be higher specifically in the cereal (*Pennisetum glaucum*) and the tuber of *Borassus aethiopum* with rates of 6.72% and 5.70 %, respectively. In addition, a positive and significant correlation was observed on the one hand between the cellulose content and that of proteins and lipids and between the protein and lipid levels. In contrast, the correlation between carbohydrate levels and those of proteins and lipids was negative and very significant. The results confirm that pulps and kernels of fruits, legumes, tubers, and cereals can be incorporated into numerous preparations intended for children to supplement their diet.

Keywords: Local food products; nutrition; food supplement; agrosylvopastoral production.

1. INTRODUCTION

In the Sahel, drought is the main risk affecting agrosylvopastoral production through reductions in food production, in the number of animals and their productivity, and in agricultural income, with the consequence of food insecurity, food crises, and nutrition [1]. Indeed, in developing countries, intake is - more monotonous in their composition, with diets lacking diversity. The degree of economic development and income level have important implications for the types of food that can be produced to satisfy human needs [2]. Thus, food prices and a country's economy are factors that influence human nutrition [3]. The Sahel is characterized by a high prevalence of malnutrition and mortality linked to undernourishment among children under 5 years of age. The causes of malnutrition are multiple factors. This can result from inadequate food intake, diseases, household food insecurity, access to health services, inappropriate care practices, access to drinking water, and poor living conditions. hygiene [4].

In Niger, malnutrition, particularly undernutrition and micronutrient deficiencies, persists and constitutes a public health problem [5]. Thus, the prevalence of global acute malnutrition (GAM) among children aged 6 to 59 months, severe acute malnutrition, and chronic malnutrition (stunting) is estimated at 12.2% (well above above the WHO safety threshold of 10%), 2.4% and 47.0% [6]. Factors associated with chronic malnutrition are age, sex, birth order, mothers' educational level, diarrhea, fever, acute respiratory infection, and anemia [7].

In addition, on a practical level, complementary feeding was introduced in a timely manner in 8 out of ten cases (80.6%), but less than half of the children (47%) received breast milk until the age of two years. The diet of children aged 6 to 23 months was generally poorly diversified (the minimum dietary diversity rate was 8.7%), mainly based on starchy foods (91.1%), followed by legumes and nuts (29.6%), fruits and vegetables rich in vitamin A (25.6%), and breast milk (85.6%). Therefore, the prevalence of overall

anemia (mild, moderate and severe) in children aged 6 to 59 months is high (55.5% [6].

The issue of nutrition is seen today as a global concern, characterized by a real collective awareness expressed within the Scaling Up Nutrition (SUN) Movement, which brought together 60 countries in 2017 [4]. One of the most promising solutions to combat malnutrition sustainably is to design programs that consider locally produced foods [8]. Dietary approaches constitute levels that can significantly contribute to reducing the prevalence of malnutrition [9]. For example, daily supplementation with moringa leaf powder significantly improves the nutritional status of children in terms of wasting, stunted growth, and underweight [9].

From the age of six months, breast milk intake is insufficient to meet the needs of infants and young children and must be supplemented by a diversified and high nutritional density so-called “complementary” diet [10]. The present study aimed to evaluate the nutritional quality of 15 plant species, from which 17 samples were collected as potential complementary foods.

2. MATERIALS AND METHODS

2.1 Presentation of the Study Area

The urban commune of Niamey (Niger) served as the study site. It comprises five (5) municipal

districts (Fig. 1). The population of Niamey is estimated to be approximately 1,492,414 inhabitants [11]. The average rainfall and temperature were 539.16 mm and 29.59°C, respectively [12]. In addition, a previous study listed 15 types of porridge used as complementary food among children aged 6 to 24 months in the urban community of Niamey, six of which come from industrial flours of known composition and the other nine are porridges. Local traditions [13]. In this region, the continuation of breastfeeding until the age of two years was estimated at 14.9%, and the minimum dietary diversity among children aged 6 – 23 months was estimated to be 31.1% [6].

2.2 Plant Material

It was made up of 15 plant species, from which 17 samples were collected. The scientific names of the plant species, organs used, vernacular names, and provenances are listed in Table 1. This table shows that six samples were used in the grilled form, eight in the raw form, and three in the cooked form.

2.3 Methodology for Determining Biochemical Parameters

The methodology used in this study varied depending on the biochemical parameters studied.




Fig. 1. Map of the study area

Table 1. Composition and characteristics of plant material

Scientific names	Names in French	Local names	Languages ¹	Organs used ²	Codes	Forms used	Origins
<i>Adansonia digitata</i>	Baobab	kuka, koo nya, bokki,	H, Z, F	Pu	Addi	believed	Local market
<i>Arachis hypogaea</i>	Peanut	gujiya, dams, kolhi,	H, Z, F	Gr	Arhy	grilled	Local market
<i>Glycine max</i>	Soy		H, Z, F	Gr	Glma	grid	Local market
<i>Ziziphus mauritiana</i>	Jujube	magaria, darey, djâbhi,	H, Z, F	Am	ZimaA	believed	Local market
<i>Western Anacardium</i>	Cashew	Say tourizé	Z	Am	Anoc	grid	Local market
<i>Borassus aethiopum</i>	Rônier	jijinia, bê, dubbhy	H, Z, F	Tu	Boae	cooked	Local market
<i>Parkia biglobosa</i>	Nere	dorowa, dosso, nareehi,	H, Z, F	Pu	Pabi	believed	Local market
<i>Cucurbita SP.</i>	Squash	kankana, laptanda, faireare.	H, Z, F	Gr	Cusp	believed	Local market
<i>vigna unguiculata</i>	Cowpea	wake, dunguri, yanpu,	H, Z, F	Gr	Viun	cooked	Local market
<i>Neocarya macrophylla</i>	cayor apple tree	gawasa, gamsa, nawudi,	H, Z, F	Pu	NemaF	believed	Local market
<i>Neocarya macrophylla</i>	cayor apple tree	gawasa, gamsa, nawudi,	H, Z, F	Am	NemaA	believed	Local market
<i>Sesamum indicum</i>	Sesame	lamti, lamti,	H, Z, F	Gr	Breast	grid	Local market
<i>Ipomoea batatas</i>	Yam	dankaly, kudaku, dagafibre,	H, Z, F	Tu	Ipba	cooked	Local market
<i>Sclerocarya birrea</i>	Marula	daniya, diney (luley), hedi,	H, Z, F	Am	Scbi	grid	Local market
<i>Pennisetum glaucum</i>	Millet	hatsi;hayni;gawuri	H, Z, F	Gr	Pegl	grid	Niamey
<i>Hyphaene thebaica</i>	Doum Palm	goriba, kangaunya, gellehy,	H, Z, F	Pu	Hyth	believed	Local market
<i>Ziziphus mauritiana</i>	Jujube	magaria, darey, djâbhi,	H, Z, F	Pu	ZimaF	believed	Local market

1: H=Hausa; Z=Zarma; F=Fulfulde. 2: Am = Almond; Pu= Pulp; Gr = Grain; Tu=Tuber.

Source: [14]

2.3.1 Water content

The method used for determining water content was proposed by AOAC [15]. It consisted of carrying out differential weighing of the crucibles, containing 5 g of sample, before and after passing through the oven at 105°C. The difference in weights before and after drying made it possible to determine the humidity, according to Equation 1.

$$W(\%) = \frac{me - ms}{me} \times 100 \quad (1)$$

W: water content in percentage (%)
me: mass of the sample before drying (g)
ms: mass of the sample after drying (g)

2.3.2 Ash content

The method used to determine ash content was described by AOAC (1990). Thus, in a crucible, 2 g of the ground material was incinerated at 550°C for 6 h in a NABERTERM brand muffle furnace, MORE THAN HEAT 30-3000°C, Germany. After cooling in a desiccator, the samples were reweighed. The ash content was expressed as mass percentage, according to Equation 2.

$$C(\%) = \frac{m2}{m1} \times 100 \quad (2)$$

m1: mass of test sample (g)
m2: mass of ash (g).

2.3.3 Cellulose

The crude fiber content was estimated using the Weende method. This method is carried out in two hydrolysis processes, one acidic with 1.25% H₂SO₄, which extracts the sugars and starch, and the other alkaline (basic) with 1.25% NaOH, allowing the proteins to be extracted and part of hemicellulose and lignin. Acid-base hydrolyses are separated and isolated by filtration and rinsing with hot water [16]. Cellulose content was calculated using the following formula, according to Equation 3:

$$\text{Cellulose brute } (\%) = \frac{P2 - P3}{P1} \times 100$$

P1: Sample weight;
P2: Weight of the crucible + cellulose + mineral materials
P3: Weight of crucible and mineral materials.

2.3.4 Protein content

The protein content of the samples was determined by the Kjeldahl method. It consists of an automatic distiller brand UDK 129. It is performed in three phases: mineralization, distillation, and colorimetric dosage (volumetric dosage). The nitrogen and crude protein contents relative to the fresh material were calculated directly and provided by the device using 6.25 as a conversion factor [17].

2.3.5 Lipid content

The fat content of the foods was determined by the Soxhlet method with gravimetric extraction of lipids. The principle is based on the extraction of a test portion with hexane by percolation. Followed by elimination of the solvent was removed by distillation, and finally by drying the residue was dried in an oven and weighed [18].

2.3.6 Total carbohydrate content

The total carbohydrate content in relation to dry matter was determined using the differential method (method used at the laboratory level). The calculation was performed using the determined values of protein, lipid, ash, and humidity [13]. The formula used was as follows (Equation 4):

$$\% \text{total carbohydrates/DM} = 100 - [\% \text{Fat} + \% \text{protein} + \% \text{ash}] \quad (4)$$

2.3.7 Determination of energy density

The energy density (ED) was calculated according to the equation of Atwater and Benedict [19].

$$DE = (9 \times \text{lipids (g)} + 4 \times \text{proteins (g)} + 4 \times \text{carbohydrates (g)} + 2 \times \text{cellulose (g)}).$$

2.3.8 Statistical analysis

The data were processed and analyzed using Minitab version 19, SPSS version 25, and Excel 13. The comparison of the means of biochemical parameters was carried out according to Tukey's test at the 5% probability threshold. Finally, the hierarchical ascending classification of foods, comparison of means according to the food category, and correlation between the different physicochemical variables were carried out using SPSS software.

3. RESULTS

3.1. Biochemical Characterization of Foods

3.1.1 Proteins, lipids, carbohydrates, ash and moisture from fruit seeds, tubers, pulps and kernels

Table 2 presents the biochemical composition of the fruit seeds, tubers, pulps, and kernels. It has a water content of less than 7%. The highest water content was 6.72% in millet seeds (*Pennisetum glaucum*) and the lowest was 1.55% (*Anacardium occidentale*). Furthermore, the total ash content was higher in the pulp of the doum palm (*Hyphaene thebaïca*) at 7.16% but lower in the seeds of millet (*Pennisetum glaucum*) at a rate of 1.85%. It is clear from this table that legumes and fruit kernels contained the highest levels of proteins and lipids. Thus, the highest protein content was observed in soybean (*Glycine max*) at 36.39%, followed by *Sclerocarya birrea* almond (30.84%), and the lowest was 2.11% in sweet potato (*Ipomoea batatas*).

Furthermore, the highest lipid content was 60.75% in *Neocarya macrophylla* kernel and 0.20% in sweet potato tubers (*Ipomoea batatas*).

In contrast, all fruit pulps, tubers, cereals, and some legumes contained sufficient carbohydrates. For this purpose, the sweet potato tuber was exclusively rich in carbohydrates at a rate of 95.25%, and the lowest value was obtained in the *Sclerocarya birrea* kernel (6.45%). The average moisture, ash, protein, lipid, and carbohydrate content was compared. These differences were not significant within each group.

3.1.2 Cellulose and energy density of seeds, pulp and kernels of fruits and tubers

Legumes and fruit kernels are extremely rich in cellulose, with the exception of *Vigna unguiculata* (Table 3). For this purpose, the highest values were reported for the almond *Neocarya macrophylla* (26.74%), the seed of *Cucurbita SP* (23.75%), almond of *Sclerocarya birrea* (18.80%), *Ziziphus mauritiana* almond (17.92%), Western *Anacardium* almond, and seeds of *Sesamum indicum* (16.29%). The cellulose content of *Vigna unguiculata* seeds is very low (1.48%). Compared to the calculated energy density, *Neocarya macrophylla* almond presented the greatest energy value. The energy values of the analyzed samples oscillate between 392.24 and 748.93 Kcal/100g.

Table 2. Protein, fat, carbohydrate, ash and moisture contents of fruit seeds, tubers, pulps and kernels

Scientific Names ¹	Proteins (100g/DM)	Fats (100g/ DM)	Carbohydrates (/100g /DM)	Ashes (100g /DM)	Humidities (100g)
<i>Adansonia Digitata</i>	3,11±0,59 ^g	0,44±0,20 ^h	90,39±2,12 ^{ab}	6,06±1,6 ^a	5,28±0,48 ^{abc}
<i>Arachis hypogaea</i>	23,76±1,01 ^c	38,93±5,68 ^d	34,01±4,88 ^f	3,3±0,27 ^{bcd}	1,69±0,51 ^g
<i>glycine max</i>	36,39±1,83 ^a	8,54±0,92 ^f	51,22±2,74 ^e	3,86±0,04 ^{bc}	3,46±0,80 ^{cdefg}
<i>Ziziphus mauritiana</i> ^A	29,40±0,43 ^b	29,79±1,04 ^e	37,79±1,09 ^f	3,02±0,13 ^{bcd}	3,85±0,74 ^{bcdef}
<i>Anacardium occidentale</i>	21,29±1,37 ^{cd}	48,92±0,40 ^c	27,60±2,12 ^g	2,19±0,48 ^d	1,55±0,77 ^g
<i>Borassus aethiopum</i>	6,18±1,50 ^f	0,83±0,40 ^{gh}	91,10±1,75 ^{ab}	1,89±0,07 ^d	5,7±0,50 ^{ab}
<i>Parkia biglobosa</i>	5,13±0,86 ^{fg}	0,89±0,24 ^{gh}	89,71±0,82 ^{ab}	4,27±0,17 ^b	3,17±0,50 ^{defg}
<i>Cucurbita SP.</i>	29,39±0,61 ^b	53,71±0,58 ^b	14,03±0,43 ^h	2,87±0,15 ^{bcd}	2,45±0,29 ^{efg}
<i>Vigna unguiculata</i>	17,82±2,35 ^d	1,94±0,25 ^{gh}	77,35±2,56 ^d	2,89±0,06 ^{bcd}	5,41±0,19 ^{ab}
<i>Neocarya Macrophylla</i> ^F	3,47±0,04 ^{fg}	0,71±0,18 ^{gh}	93,01±1,13 ^{ab}	2,81±0,91 ^{bcd}	5,58±0,99 ^{bcde}
<i>Neocarya Macrophylla</i> ^A	18,89±1,48 ^d	60,75±0,45 ^a	18,29±2,24 ^h	2,07±0,38 ^d	2,19±0,22 ^{fg}
<i>Sesamum indicum</i>	24,01±0,66 ^c	60,05±0,53 ^a	12,96±0,18 ^f	2,98±0,22 ^{bcd}	1,86±0,30 ^g
<i>Ipomoea batatas</i>	2,11±1,11 ^g	0,20±0,10 ^h	95,25±1,36 ^a	2,44±0,64 ^{cd}	3,17±0,58 ^{defg}
<i>Sclerocarya birrea</i>	30,84±1,86 ^a	58,64±1,54 ^a	6,45±1,28 ⁱ	4,07±0,03 ^b	1,76±0,52 ^g
<i>Pennisetum glaucum</i>	10,06±0,90 ^e	5,17±0,41 ^{fg}	82,92±1,60 ^{cd}	1,85±0,38 ^d	6,72±0,43 ^a
<i>Hyphaene thebaïca</i>	3,66±1,41 ^g	0,78±0,14 ^{gh}	88,41±1,65 ^{bc}	7,16±0,15 ^a	4,43±0,85 ^{bcd}
<i>Ziziphus mauritiana</i> ^F	4,15±0,40 ^g	0,85±0,39 ^{gh}	90,87±0,71 ^{ab}	4,12±0,34 ^b	4,17±1,15 ^{ab}

1: F=Fruit Pulp and A=Fruit Kernel. The means on a column sharing no letters are significantly different at the 5% threshold

Table 3. Water, ash, cellulose and energy density contents

Scientific Names ¹	Cellulose (g /100g /DM)	Energy density (Kcal/100g)
<i>Adansonia Digitata</i>	8.10±0.23	394.13
<i>Arachis hypogaea</i>	11.35±0.20	604.13
<i>glycine max</i>	7.84±0.60	442.95
<i>Ziziphus mauritiana</i> ^A	17.92±1.29	572.67
<i>Western Anacardium</i>	16.42±3.48	668.69
<i>Borassus aethiopum</i>	2.82±0.24	402.21
<i>Parkia biglobosa</i>	14.08±1.28	415.52
<i>Cucurbita SP.</i>	23.75±2.19	704.54
<i>Vigna unguiculata</i>	1.48±0.33	401.06
<i>Neocarya Macrophylla</i> ^F	4.68±0.10	401.65
<i>Neocarya Macrophylla</i> ^A	26.74±1.75	748.93
<i>Sesamum indicum</i>	16.29±2.12	720.91
<i>Ipomoea batatas</i>	2.61±0.11	396.43
<i>Sclerocarya birrea</i>	18.80±3.31	714.50
<i>Pennisetum glaucum</i>	2.76±0.25	423.99
<i>Hyphaene thebaica</i>	10.46±0.23	396.18
<i>Ziziphus mauritiana</i> ^F	2.23±0.10	392.24

1: F=Fruit Pulp and A=Fruit Kernel

Table 4. Comparison of nutrient content based on food category

Food categories	Proteins	lipids	Carbohydrates	Humidities	Ashes	Cellulose
Pulps	3.90	0.73	90.48	4.53	4.88	7.91
Cereals	10.06	5.17	82.92	6.72	1.85	2.76
Legumes	26.27	32.63	37.91	2.97	3.18	12.14
Almonds	25.11	49.52	22.53	2.34	2.84	19.97
Tubers	4.14	0.51	93.18	4.44	2.17	2.72
Averages	15.86	21.83	58.90	3.67	3.40	11.08

3.1.3 Correlation according to biochemical parameters of fruit seeds, tubers, pulps and kernels

The results of comparisons of the averages of biochemical parameters according to the food category to which the plant species belongs (Table 4), showed that legumes are rich in proteins and they are followed by fruit kernels. Fruit almonds had the highest lipid content. They were also the most abundant in cellulose. Furthermore, the highest carbohydrate content was observed in the pulp, cereal, and tuber categories. Finally, the vast majority of mineral elements were found in fruit pulps.

The ascending hierarchical classification made it possible to identify three different groups:

-Group 1: Composed of foods such as the tubers of the young shoots of *Borassus aethiopum* (Boae) and *Ipomoea batatas* (Ipba), the fruit pulps of *Ziziphus mauritiana* (ZimaF),

Neocarya macrophylla (NemaF), *Adansonia digitata* (Addi), *Hyphaene thebaica* (Hyth) and *Parkia biglobosa* (Pabi), the seeds of *vigna unguiculata* (Viun) and *Pennisetum glaucum* (Pegl). It represents 53% of foods and is characterized by high values of carbohydrates and water. However, these foods have low values of proteins, lipids and celluloses.

- Group 2: Composed of foods such as *Sesamum indicum* (Breast) legumes and *Cucurbita SP* (Cusp), the fruit kernels of *Sclerocarya birrea* (Scbi) and *Neocarya macrophylla* (NemaA). It represents 29% of the samples and is characterized by high protein values.

-Group 3: Composed of foods such as *Arachis hypogaea* (Arhy) and *Western Anacardium* (Anoc) and *Ziziphus mauritiana* (ZimaA). It represents 18% and is characterized by high values of lipids and cellulose but low values of carbohydrates and water.




Fig. 2. Ascending hierarchical classification of samples

Table 5. Correlation between the different physicochemical variables

		Humidity	Ash	Cellulose	Protein	Lipid	Carbo
Humidity	Corrélacion	1					
	Sig. (bilatérale)						
Ash	Corrélacion d	0,053	1				
	Sig. (bilatérale)	0,710					
Cellulose	Corrélacion	-0,683**	-0,034	1			
	Sig. (bilatérale)	0,000	0,815				
Protein	Corrélacion	-0,542**	-0,214	0,565**	1		
	Sig. (bilatérale)	0,000	0,132	0,000			
Lipid	Corrélacion	-0,751**	-0,278*	0,837**	0,693**	1	
	Sig. (bilatérale)	0,000	0,048	0,000	0,000		
Carbo ¹	Corrélacion	0,741**	0,236	-0,814**	-0,848**	-0,968**	1
	Sig. (bilatérale)	0,000	0,095	0,000	0,000	0,000	

** The correlation is significant at the 0.01 level (two-tailed). * The correlation is significant at the 0.05 level (two-tailed). ¹ Carbohydrate.

Table 5 presents the various correlations established between the biochemical measurements. The biochemical parameters from the samples showed positive and significant correlations ($r = 0.56$, $P = 0.0000$; $r = 0.83$, $P = 0.0000$ and $r = 0.69$) respectively between the cellulose content and the protein and lipid levels, then between the protein level and the lipid level. On the other hand, a negative and very significant correlation ($r = -0.84$, $P = 0.0000$ and $r = -0.96$; $P = 0.0000$) was observed between the level of carbohydrates and those of proteins and lipids. Likewise, negative and significant correlations ($r = -0.75$, $P = 0.0000$; $r = -0.68$, $P = 0.0000$; $r = -0.54$, $P = 0.0000$) were observed between the humidity level and the lipid level, between the humidity level and the cellulose level as well as between the protein level and the humidity level.

4. DISCUSSION

Niger has plant species that provide numerous products that are used by the general population, particularly in rural areas. The food use of the latter mainly concerns fruits, pulps, leaves, flowers, seeds, and almonds [20]. The present study involved 15 local plant species, of which 17 were distributed as follows: 4 fruit pulps, 4 fruit almonds, 6 seeds, 2 tubers. Biochemical analyses of all these samples showed very variable contents of proteins, lipids, carbohydrates, cellulose, moisture, and ash (mineral elements), which often vary depending on the plant species, food group, and part concerned (fruit or almond). It is important to note that all data were reported on a dry-matter basis. Most of the previous studies encountered were more interested in raw fibers (total fibers) than cellulose, as well as the different types of minerals and ash contents; hence, these points will be less compared with literature data.

4.1 Moisture, Ash, Proteins, Lipids and Carbohydrates

Biochemical characterization of fruit seeds, tubers, pulps, and kernels showed water contents lower than 7%. This low content allows for better conservation [21]. These rates are lower than those reported for Togo on seeds (9%) [8]. In the present study, the highest water content was 6.72% for millet seeds (*Pennisetum glaucum*). It was 5.28% in the pulp of *Adansonia digitata*, 2.45% in the seeds of *Cucurbita SP*, and 1.55% in the kernel of *Western Anacardium*,

which was the lowest value. These values differ from those of another study conducted in Niger on two species of Cucurbitaceae (*Citrullus Colocynthis* and *Lagenaria siceraria*), whose water content varied from 7 to 9.50% [22]. The baobab pulp was characterized by a water content that was close to that found by other authors, which varied from 6 to 15% [23]. In addition, baobab pulp is characterized by its prebiotic potential. Indeed, the fermentation of baobab pulp powder presents characteristics of selective utilization by host microorganisms [24]. The highest total ash contents were observed in the pulps of *Hyphaene thebaïca*, *Adansonia digitata*, *Parkia biglobosa*, *Ziziphus mauritiana* with respective values of 7.16%, 6.06%, 4.27%, 4.12% and that of the lowest in millet seeds (*Pennisetum glaucum*) of 1.85%. A high total ash content of 9.7% has been reported for baobab pulp [25].

The present study shows that legumes and fruit kernels contain the highest levels of proteins and lipids. The highest protein content was observed in soybean (*Glycine max*) at 36.39%, followed by *Sclerocarya birrea* almond (30.84%), and the lowest was 2.11% in sweet potato (*Ipomoea batatas*). The results obtained are approximately similar to those found in soy varieties consumed in Burkina Fasso and Togo, with protein contents of 31.04% and 34.43%, respectively [8,26]. According to a previous study in Niger, the kernels of *S. birrea* fruits constitute a great potential and a very rich composition, reporting contents that vary depending on the origin from 31.1 to 39.9% of proteins [27]. It is a very fruitful plant, the number of fruits per tree varies from 136 to 4256 with an average fruit weight and diameter of 12.66 and 26.71 mm respectively and the number of almonds varies from 1 to 3 per fruit [28].

The highest lipid content was obtained in the kernels of *Neocarya macrophylla*, *Sclerocarya birrea*, and *Anacardium ouest*, the seeds of *Sesamum indicum* and *Cucurbita*, respectively 60.75%, 58.64%, 48.92%, 60.05% and 53.71%. These values were higher than those of the *Ipomoea tuber batatas* and fruit pulps of *Adansonia digitata*, which were 0.20% and 0.44%, respectively. A previous study by Niger noted a higher fat content of 67.5% for the almond *Neocarya macrophylla* [29]. The observed variations could be linked to the degree of maturity, genetic influence, temperature, sunshine, and nature of the soil. The results of another previous study on the almond of this

plant showed that the oil contained myristic acid, palmitic acid, stearic acid, palmitoleic acid, elaidic acid, oleic acid, erucic acid, behenic acid, heneicosanoic acid, icosatetraenoate and eicosatrienoic acid [30,31]. Considering its nutritional benefits, this plant species must be subject to sustainable management because *N. macrophylla* is mainly renewed by sowing (87.6%) and only slightly by stump shoots (11.1%) [32]. Work carried out on *Sclerocarya birrea* almonds in Niger reported contents that varied depending on the origin, from 38 to 56.3% fat, 4.58 to 22.5% carbohydrates, 3.60 to 4, and 58% ash [27]. Other studies have shown variability in nutritional characteristics among the provenances of *Western Anacardium*. Thus, the highest fat values were noted in nuts from Ziguinchor (54%), followed by those from Sédhiou (44%) and Kolda (37%) [33]. In addition, studies have found similar lipid levels in *Sesamum indicum* of 60.05%, 61.70%, 52.4 to 62.8%, and 49.83% to 59.85%, respectively [34–36].

The present study also showed that all fruit pulps, tubers, cereals, and some legumes contained sufficient carbohydrates. Indeed, the tubers of sweet potato (*Ipomoea batatas*) and the young shoot of *Borassus aethiopum*, the fruit pulps of *Neocarya macrophylla*, *Ziziphus mauritiana*, and *Adansonia digitata* are exclusively rich in carbohydrates with respective levels of 95.25%, 91.1%, 93.01%, 90.87%, and 90.39%, and the lowest value was obtained in the kernel of *Sclerocarya birrea* (6.45%). In Burkina Faso, a similar study on the tuber of *Borassus aethiopum* found a similar carbohydrate content of 86.75% [37]. However, different results were reported for the fruit pulp of *Ziziphus mauritiana*, which was 13.05% [38]. This difference could be explained by the methodology used to determine carbohydrate content (spectrophotometric assay). Similar to *Adansonia digitata* fruit pulp, different results were reported by Benin 73.12% [39]. This difference in carbohydrate content could be due to the differential method used during the operation, as it inserted the cellulose content.

4.2 Celluloses and Energy Density

As shown in Table 3, legumes and fruit kernels are extremely rich in cellulose, with exception of *Vigna unguiculata*. Thus, the highest values were reported for the *Neocarya macrophylla* kernel (26.74%), the seeds of *Cucurbita SP* (23.75%),

almond of *Sclerocarya birrea* (18.80%), almond of *Ziziphus mauritiana* (17.92%), almond of *Anacardium occidentale*, seed of *Sesamum indicum* (16.29%), and seed of *Vigna unguiculata* are very poor in cellulose (1.48%). Previous studies have reported different cellulose results for *Western Anacardium*, which varied according to provenance, including Kolda (14%), Ziguinchor (13.5%), and Sédhiou (9%) [33].

Compared to the calculated energy density, *Neocarya macrophylla* almond had the highest energy value. The energy values of the analyzed samples oscillate between 392.24 and 748.93 Kcal/100g. The energy richness of fruit almonds and certain legumes could be linked to their high fat content. [8] reported energy values lower than those of this study in the seeds of two legumes *Arachis hypogaea* and *Sesamum indicum* of 576.08 Kcal and 500.29 Kcal respectively. This difference could be explained by climatic factors and differences in the methodology used to determine carbohydrate levels.

4.3 Correlation according to Biochemical and Physicochemical Parameters of Seeds, Tubers, Pulps and Fruit Kernels

The results of the comparison of the averages of the biochemical parameters according to the food category to which the plant species belongs (Table 5) show that legumes have the highest average protein content of 26.27%, which is very close to that of fruit almonds (25.49 %). This could mean that regardless of whether legumes are available, fruit kernels can be used for their important role in the development, maintenance, and repair of tissues. In addition, fruit almonds had the highest lipid content (49.52 %), followed by that of legumes (32.63 %). However, the fat content varies, depending not only on the species, but also on the part of the fruit studied (pulp or kernel) [40].

In addition, the highest carbohydrate values were observed in the pulp, cereal, and tuber categories with rates of 90.48%, 82.92% and 93.18%, respectively. Almonds were the richest in cellulose (19.97 %), followed by legumes (12.14 %). As for mineral elements, the largest majority was found in fruit pulps with a content of 4.88%, followed by legumes come in second position with a value of 3.18%. The highest water content was observed in cereals (6.72%), followed by pulp (4.53%), which is practically the

same for tubers. It emerges from the analysis of this table that all categories of food rich in proteins and lipids have an inverse relationship with the quantity of water. Indeed, foods with the greatest quantities of proteins and lipids were the poorest in water. These results are different from those found in research work in Algeria, where all the energy-rich fruits studied were in an inverse relationship with the quantity of water, and fruits with a high energy content had the poorest water content [40]. This observation is statically confirmed in Table 5 with negative and significant correlations at the 0.01 level ($r = -0.75$, $p=0.0000$; $r = -0.68$, $p= 0.0000$; $r= - 0.54$, $p=0000$), which were observed between the humidity level and the lipid level, as well as between the humidity level and the cellulose level, as well as between the protein and lipid level.

Hierarchical ascending classification of the samples revealed three large groups according to their nutritional composition (Figure 2). The first group comprised 53% of the samples and mainly consisted of fruit pulps and starchy foods (tubers) (Boae, ZimaF, NemaF, Ipba, Addi, Hyth, Pabi, Viin, and Pegl). The discriminating criterion was the richness of fruit pulps, starches, and *Vigna unguiculata* in carbohydrates and water. The second group constituted 29%, namely Seal Scbi, Cusp, and Nema. The characteristic criterion of this group was a high protein value. The third group included 18% of the samples studied, namely Arhy, Anoc, and ZimaA. It is mainly characterized by high values of lipids and cellulose but low values of carbohydrates and water.

5. CONCLUSION

At the end of this work, we noted that the plant species studied presented remarkable potential from a nutritional point of view. Plant proteins are present in greater quantities in all almonds and legumes (36.39% for *glycine max*) and can be used in different areas such as human food and livestock feed. The oils of fruit kernels and oilseed legumes are present in appreciable quantities, including 60.75% in the *Neocarya macrophylla kernel*, and can have food uses. Pulps, tubers, and cereals exhibited significant energy contributions.

All parts of the plants analyzed are already used in food in Niger, and it is possible to consider different transformations as flour and incorporate it into different food products, either as food

supplements or as raw materials.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Moussa BMC, Bio MT, Halima OD, Abdourahamane B. Climate risks and food and nutritional security in Niger: mapping of impacts and resilience needs. *Journal in Environmental Sciences*.2022;22(1):1492-8442.
2. Westphal E, Embrechts J. Tropical food crops: with special reference to Cameroon; Communication; Pudoc: Wageningen. 1985;(4). ISBN 978-90-220-0845.
3. Djibo DH. Eating practice of mothers in Niamey: between rule and reality, what future for children. *Review of Multidisciplinary Studies in Economic and Social Sciences*.2017;(6):2489-2068.
4. UNICEF National Strategy for the prevention of different forms of malnutrition in Djibouti. 2022;170.
5. Alirou MA, Mamadou L, Chaibou Y, Aboubacar A, Mahamadou A, Aboubacar M, Fatimata JA and al. Study of the frequency of consumption of food recipes at the community level in five regions of niger. *International Journal of Current Research*.2023;15(2):23784-23795.
6. National Institute of Statistics (INS). Retrospective Nutritional and Mortality Survey in Niger. 2022;96.
7. Ali LZ, Issa AA, Attinine AS, Daouda S, Salissou LB, Mamadou ML. Factors associated with chronic malnutrition among children aged 0-59 months in Niger. *Health Research in Africa*. 2023;1(3):54-59.
8. Souho T, Bomboma D, Compaoré EWR, Essowèréou AA, Ouadja B, Kabassina M, Pilo M, Epse Yao RRA, Amouzou KS. Grains production in high climate change impacted regions and its potential for the supply of critical nutrients for humans nutritional wellbeing. *Food Production, Processing and Nutrition*.2023;5(31):2-12.
9. Houndji BVS, Bodjrenou SF, Londji SBM, Ouetchehou RA, Acakpo A, Amouzou K,Hounmenou D. Improvement of the nutritional status of children aged 6 to 30 months in Lissèzoun (Central Benin) by

- Moringa Oleifera (Lam.) leaf powder. International Journal of Biological and Chemical Sciences. 2013;7(1):225–235.
10. Olive F, Mouquet-Rivier C, Nelly F, Bichard A, Boulle-Martinaud N, Kaboré C, Denizeau M, Zagre N, Dain A, Ndiaye N, et al. The Locally Produced Infant Flour Sector in 6 Sahelian Countries. 2020;164.
 11. National Institute of Statistics (INS). Distribution of the Population of the Niamey Region by Age Group According to Sex in 2024. 2024;103.
 12. Boubacar G, Saley MM, Saley IA, Mouhaimouni M, Fodé M. Statistical analyzes of variations in temperatures and precipitation observed in Niamey and Nguigmi. Meteorology. 2020;103:28–33.
 13. Garba dit Gado Halidou O, Lewamy M, Chaibou C, Adamou OA, Sabo HS. Inventory and Nutritional analysis of porridges used as complementary food in children aged 6 to 24 months in the urban community of niamey, Niger. EWASH & TI Journal. Environmental and Water Sciences, Public Health & Territorial Intelligence. 2019;3(2):2-8.
 14. Inran. Illustrated Glossary of Some Plants OF Niger. 2017;52.
 15. AOAC. Official methods of analysis. 15th Edition, A.o. Washington DC: OAC; 1990.
 16. Chehma A, Faye B, Bastianelli D. Nutritional Values of Perennial Plants from Algerian Saharan Rangelands for Camels. 2010;204:263–268.
 17. AOAC Official method of analysis of the association of official analytical chemists. Ed AOAC. Washington DC. 1984;P24003.
 18. AOAC Official Methods of Analysis (16th Edn). Association of Official Analytical Chemists. 1995;771.
 19. Atwater WO BF. Experiments on the metabolism of energy in the human body, 1898-1900. United States. Office of Experiment Stations bulletin N°109. Washington; 1902.
 20. Dan Guimbo I, Barage M, Douma S. Preliminary studies on the food use of spontaneous plants in the peripheral areas of the W park of Niger. International Journal of Biological and Chemical Sciences. 2013;6(6):4007–4017.
 21. Donatien A, Martial G, Queen B, Such KR, Welcome C, Pascal A, Eugenie A. Biochemical profile and antioxidant activity of parkia biglobosa and tamarindus indica fruits acclimated in benin. International Journal of Advanced Research. 2018;6(11):702-711.
 22. Sabo HS, Sadou H, Amoukou I, Alma M, Mahamane S. Potentials in Vegetable oils and various nutrients of eighteen varieties of Niger Cucurbitaceae. International Journal of Biological and Chemical Sciences. 2014;8(3):1075-1088.
 23. Cisse M, Sakho M, Dornier M, Diop CM, Reynes M, Sock O. Characterization of the baobab fruit and study of its transformation into nectar. Fruits, Cirad/EDP Sciences. 2009;64:19–34.
 24. Foltz M, Zahradnik AC, Van den AP, Ghyselinck J, Marzorati M. A Pectin-Rich, Baobab Fruit Pulp Powder Exerts Prebiotic Potential on the Human Gut Microbiome in vitro. 2021;9:2-14.
 25. AYOSSO JOG. Diversity and nutritional value of local food resources for the nutrition of children 6-24 months. Faculty of agricultural sciences, University of Abomey-Calavi, Master Memory. 2015;90.
 26. Hama-Ba F, Siedogo M, Ouedraogo M, Dao A, Dicko H, Diawara B. Consumption patterns and nutritional value of food legumes in Burkina Faso. African Journal of Food, Agriculture, Nutrition and Development. 2017;17(4):12871-12888
 27. Arohalassi HM, Gounga M, Issa AR. Biochemical characterization of the pulp and kernel of ripe fruits of Sclerocarya birrea (A. Rich.) Hochst. consumed in Niger. International Journal of Biological and Chemical Sciences. 2022;16(6):2560-2572.
 28. Bationo-Kando P, Sawadogo B, Kiebre Z, Kientega P, Sawadogo N, Nanema K, Traore E, Sawadogo M, Zongo J. Productivity Characteristics and Development Strategies of *Sclerocarya birrea* in Burkina Faso. African Crop Science Journal. 2016;24(1):35 - 47.
 29. Balla A, Baragé M, Larwanou M, Adam T. Endogenous Know-How in the Food valorization of the fruits of the cayor apple tree (*Neocarya Macrophylla*) in Niger. Bulletin of Agronomic Research of Benin. 2008;59:1–8.
 30. Niamke A, Assanvo J, Taloute R, Allico D. Study of the physico-chemical, functional, granulometric, mineral and antioxidant properties of three (3) Flours FS, F40, F50 of young shoots of roan (*Borassus Aethiopum* Mart.) According to the drying method. Journal of Experimental Agriculture International. 2022;44(10):15-28.

31. Warra AA, Hassan LG, Babatola L J, Omodolapo AA, Ukpanukpong UK, Berena GA. Characterization of neocarya macrophylla seed Oil using gas chromatography-mass spectrometry (GC-MS) and fourier transform infra-red (FT-IR) Techniques. Chemical Science International Journal. 2019;28 (3):1-7.
32. Aboubacar K, Douma S, Mamoudou BM, Seyni RSD. Structure of natural populations of neocarya macrophylla (sabine) prance, woody of food interest, in Dallol Bosso, Niger. Tropical wood & forests .2018;337(3):67–78.
33. Samb CO, Biteye M, Faye E, Diarra R, Diaw N, Thiam M, Fall B. Agromorphological, chemical and biochemical characterization of raw nuts from three cashew trees (Anacardium Oest l.) From southern senegal of cashew tree. African Agronomy. 2021; 33(2):177 - 189.
34. El-Beltagi HS, Maraai RW, El-Ansary AE, Rezk AA, Mansour AT, Aly AA. Characterizing the bioactive ingredients in sesame oil affected by multiple roasting methods. Food .2022;11:1-21.
35. Makinde FM, Akinoso R. Comparison between the nutritional quality of flour obtained from raw, roasted and fermented sesame (*Sesamum indicum* L.) seed grown in Nigeria. Acta Sci Pol Technol Food. 2014;13(3):309-319.
36. Wei P, Zhao F, Wang Z, Wang Q, Chai X, Hou G, Meng Q. Sesame (*Sesamum indicum* L.): A comprehensive review of nutritional value, phytochemical composition, health benefits, development of food, and industrial applications. Nutrients. 2022;14:1-26.
37. Zongo O, Nikiéma NW, Kaboré S, Hama C, Sawadogo A, Zoungrana BA, Nikiéma F, Traoré Y, Savadogo A. Evaluation of nutritional composition of borassus aethiopum mart. hypocotyl (Koboula) consumed in Burkina Faso. American Journal of Food Science and Technology. 2019;7(6):215-222.
38. Makalao MM, Savadogo A, Zongo C, Traore AS. Nutritional composition of 10 wild fruits consumed in three departments of Chad. International Journal of Biological and Chemical Sciences. 2015;9(5):2385-2400.
39. Fassinou FTK, Chadare FJ, Affaffere M, Gbaguidi MA. Mutchayan a derived product from baobab fruit pulp used as complementary food in Benin. Research Application Summary. 2019;18 (1):46 - 53.
40. Abdeddaim M. Study of the biochemical composition of the fruits of five plant species present in the Aurès region with a view to their food or pharmacological use. (*Celtis australis* L, *Crataegus azarolus* L, *Crataegus monogyna* J, *Elaeagnus angustifolia* L, and *Zizyphus lotus* L). Doctorate in Sciences, ferhat abbas sétif 1 University: Faculty of Natural and Life Sciences. 2016;174.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/116415>