



Spatial and Temporal Trend Analysis of Precipitation and Temperature Data in Malwa Region of Kshipra River Basin, India

Pramod Kumar Meena ^{a++*}, Deepak Khare ^{b#},
Surendra Kumar Chandniha ^{c†} and Harithalekshmi V ^{d‡}

^a Department of Agricultural Engineering, Betul, Ministry of Agriculture and Farmers Welfare, Madhya Pradesh, India.

^b Indian Institute of Technology, Roorkee, Uttarakhand, India.

^c BRSM CAET and RS, IGKV, Mungeli, India.

^d Department of Agrometeorology, College of Agriculture, IGKV, Raipur, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author PKM did the investigation, methodology, coding, scripting, mapping and GIS analysis. DK did the project administration and supervision. Author SKC did the software data curation and formal analysis. Author HV writing review and editing. All authors writing original draft. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijecc/2024/v14i64212>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/118265>

Original Research Article

Received: 28/03/2024

Accepted: 02/06/2024

Published: 06/06/2024

⁺⁺ Assistant Engineer;

[#] Professor;

[†] Assistant Professor and Scientist;

[‡] Research Scholar;

*Corresponding author: E-mail: pramodcae@gmail.com;

Cite as: Meena, Pramod Kumar, Deepak Khare, Surendra Kumar Chandniha, and Harithalekshmi V. 2024. "Spatial and Temporal Trend Analysis of Precipitation and Temperature Data in Malwa Region of Kshipra River Basin, India". *International Journal of Environment and Climate Change* 14 (6):67-85. <https://doi.org/10.9734/ijecc/2024/v14i64212>.

ABSTRACT

This study investigated trends in climatic variables, such as rainfall and temperature (mean, maximum, and minimum), on both annual and seasonal basis (pre-monsoon, monsoon, post-monsoon, winter) at four stations viz. Mahidpur, Alot, Dewas, and Indore in the Kshipra River basin, India. Using the non-parametric Mann-Kendall test and Sen's slope estimator, trends were analyzed for data from 1980 to 2012. The analysis revealed statistically significant positive trends in maximum, minimum, and average temperatures on both annual and winter scales across all stations. Additionally, a significant positive trend in average temperature was observed in the post-monsoon season, except at Alot station. The annual trend magnitude for average temperature (T_{avg}) ranged from $0.020^{\circ}\text{C}/\text{year}$ (Indore) to $0.024^{\circ}\text{C}/\text{year}$ (Mahidpur). Rainfall trends on an annual scale showed a non-significant increase at Indore and Alot, while Mahidpur and Dewas exhibited a decreasing trend. Over the 32 year period, 2000 and 1993 were identified as breakpoint years for rainfall and temperature, respectively. From 1993 to 2012, annual minimum, maximum, and average temperatures increased by approximately 1.5%. Findings of this study can be used as a valuable information for water resource and crop planning, policy making and preparation of contingency plans.

Keywords: Trend; mann-kendall; spatial; temperature; rainfall; Kshipra; basin; India.

1. INTRODUCTION

The changing climatic conditions and warming of environment has major impact on the water resources and agricultural sectors, and overall economy of the nation. Climate change may increase the frequency and intensity of extreme precipitation and temperature events; these impacts can be explored through trend analysis [1]. Due to intensification of hydrological cycle, drastic and oppressive change in hydrologic parameters (like precipitation, evaporation and stream flow) have occurred which influences significantly the flow regimes. Among various hydrologic parameters which govern the hydrological processes, temperature has a significant and direct effect on nearly all such variables [2]. Air temperature is generally recognised as a good indicator of the state of climate globally because of its ability to represent the energy exchange process over the earth's surface with reasonable accuracy [3,4]. However, the high temperature change warms the climate which leads to intensification of hydrological cycle such as higher rates of evaporation and increase of liquid precipitation. Some studies on temperature variation on global scale [5,6,7] have established the fact that the earth's atmosphere has witnessed a significant warming in last century. The studies in mountainous areas like the Swiss and Polish Alps, the Rockies [8,9,10,11,12] and the Andes [13] have demonstrated significant rise in air temperatures with alarming effects on their environment.

Several studies in India had also been carried out to determine the changes in temperature and rainfall along with its association with climate change. A study by [14] in the north-western Himalaya region showed a significant rise in air temperature, with winter warming occurring at a faster rate. [15] found the increasing temperature and decreasing precipitation trend at northern Himalaya region. Similarly [16] also found an increase of 0.9°C in annual maximum temperature over the western Himalaya. In another study conducted by [17] demonstrated the variability and trends using annual and seasonal temperature variables for the 45 station in central India. They found a significant increase of $0.62^{\circ}\text{C}/102$ years in the annual minimum temperature, and $0.60^{\circ}\text{C}/102$ years in the annual maximum and mean temperature. A trend analysis of mean monsoon rainfall and mean summer temperature conducted by [18] in Karnataka, Gujarat, Rajasthan, and Maharashtra revealed decreasing rainfall trends in all states except Maharashtra, which had a slightly negative trend. Temperature trends increased in all states except Maharashtra, where the annual temperature trend was positive.

The changes in temperature, precipitation and other related climatic variables are likely to influence the amount and distribution of runoff in all river systems globally [19,20,21,22]. The non-parametric methods such as Mann-Kendall trend test, Spearman's tau, Sen's slope, regression line, and Sen's innovative trend analysis are widely used for analysing the trends in a number of hydro-climatic time series namely rainfall,

temperature, pan-evaporation, wind speed etc. [23,24,25,26,27,28,29,30,31,32,33,34,35,36]. No detailed study of trends for climatic data, in particular temperature and precipitation for the Malwa region of Kshipra river basin has been reported. Keeping in the view, the present study aims is to detect and quantify the trends of climatic variables such as precipitation and Temperature by using non-parametric statistical tools. The Mann-Kendall test and Sen's estimator of slope method (non-parametric) have been applied to determine the trend of temperature and rainfall series available for various meteorological stations distributed over the Kshipra river basin. The trends and their magnitude for these climatic variables have been determined for both annual as well as seasonal time scale. In addition, the shift analysis has also been performed using Pettitt's method.

2. MATERIALS AND METHODS

2.1 Study Area

The study catchment Kshipra is located in the western part of Madhya Pradesh state, India. It falls in the semi-arid sub-tropical type climatic region of India (Fig. 1). Geographically, the Kshipra watershed is located between 75° 25' 04'' to 76° 13' 19'' E longitude and 22° 27' 29'' to 23° 56' 40'' N latitude. It covers a geographical area of 5617 km² having elevation ranges from 444 m at the outlet and 819 m at the ridge line in upper reaches. The slope of the watershed is upto 15%. Approximately 51% of the area is occupied by agriculture followed by fallow land, barren land and open forests. The built-up area can be found only at around Indore, Ujjain and Dewas cities. The soil of the watershed is classified as Clay (38.28%), Loam (31.72%), Sandy loam (0.07%) and Silt clay (29.85%) (NBSSLUP, Nagpur). The annual rainfall ranges between 581.94 mm and 1299.13 mm, and most (about 80%) of the rainfall is concentrated in the months July, August and September (or known as monsoon season in India). Daily mean temperature ranges from a maximum of 45.5 °C (May) to minimum of 6 °C (January). The water

of the Kshipra River is being used for drinking, industrial and irrigation purposes. Apart from this, the river also has spiritual importance and treated as holy and sacred. Ujjain one of the ancient and shrine city is located on the bank of Kshipra River, where Maha-Kumbh Mela (Hindus big pilgrimage fair) organized once in every 12 years.

2.2 Meteorological Data Used

The meteorological data such as temperature and rainfall for the stations Dewas, Indore, Mahidpur and Alot was collected from office of the State Data Centre situated at Bhopal, Madhya Pradesh, India. The detail of characteristics of each station is given in Table 1. A total of 32 years (1980-2012) of daily temperature (mean, maximum and minimum) and rainfall data have been used for the trend analysis.

2.3 Data Processing

The long term (1980-2012) data for the temperature and rainfall were compiled and processed for the seasonal and annual trends analyses. The available daily temperature data series has been used to compute the monthly time series as maximum (T_{max}), minimum (T_{min}), average (T_{avg}), for all the meteorological stations. The daily rainfall data series of four stations namely Dewas, Indore, Alot and Mahidpur were used to form monthly totals. To investigate the changes in climatic variables at different time scales, a year was divided into four principal seasons: Pre-monsoon season (March to May), Monsoon season (June to September), Post-monsoon season (October to November) and Winter season prevailing (December to February).

The monthly data of temperature and rainfall were further used to compute the annual and seasonal time series, which were in turn used for the investigation of trend on seasonal and annual time scale.

Table 1. Details of the meteorological stations located in the study area

Sl. No.	Station	Latitude	Longitude	Altitude (m)
1	Dewas	22° 58' 45"	76° 04' 00"	535
2	Indore	22° 43' 00"	75° 54' 30"	563
3	Mahidpur	23° 28' 30"	75° 39' 20"	463
4	Alot	23°45' 55"	75° 33' 25"	450




Fig. 1. Location map of Kshipra river basin

2.4 Methodology

In the present study, to analyze the trends of the climatic series of Temperature and Rainfall for each individual station, the popular statistical non-parametric methods such as Mann-Kendall test and Sen's estimator of slope method have been used. The two phase systematic approach has been adopted to determine the trend. Firstly, the Mann-Kendall test was used for the presence of a monotonic increasing or decreasing trend in the time series. Secondly, the Sen's estimator of slope test was used to determine the magnitude of the trend in the time series. The details of both the methods are described in the following sections.

2.4.1 Sen's estimator of slope

The magnitude of trend in a time series was determined using a non-parametric method known as Sen's estimator [37]. This method assumes a linear trend in the time series and has been widely used for determining the magnitude

of trend in hydro-meteorological time series [38,39,40]. In this method, the slopes (T_i) of all data pairs are first calculated as:

$$T_i = \frac{x_j - x_k}{j - k} \text{ for } i = 1, 2, \dots, N \quad (1)$$

Where x_j and x_k are data values at time j and k ($j > k$), respectively. The median of these N values of T_i is Sen's estimator of slope which is calculated as:

$$\beta = \begin{cases} T_{\frac{N+1}{2}} & \text{if } N \text{ is odd,} \\ \frac{1}{2} \left(T_{\frac{N}{2}} + T_{\frac{N+2}{2}} \right) & \text{if } N \text{ is even.} \end{cases} \quad (2)$$

A positive value of β indicates an upwards (or increasing) trend and a negative value indicates a downwards (or decreasing) trend in the time series.

2.4.2 Mann–Kendall test

To ascertain the presence of a statistically significant trend in hydrologic climatic variables such as temperature and precipitation with reference to climate change, the non-parametric Mann–Kendall (MK) test has been employed by a number of researchers [41,42,43,44,45]. The MK method searches for a trend in a time series without specifying whether the trend is linear or non-linear. It checks the null hypothesis of no trend versus the alternative hypothesis of the existence of an increasing or decreasing trend. The statistic S for MK test is defined as:

$$S = \sum_{i=1}^{N-1} \sum_{j=i+1}^N \text{sgn}(x_j - x_i) \quad (3)$$

Where, sgn is sigma function; the x_j are the sequential data value, N is the number of data points. Assuming $(x_j - x_i) = \theta$, the value of $\text{sgn}(\theta)$ is computed as follows:

$$\text{sgn}(\theta) = \begin{cases} 1 & \text{if } \theta > 0, \\ 0 & \text{if } \theta = 0, \\ -1 & \text{if } \theta < 0. \end{cases} \quad (4)$$

This statistic represents the number of positive differences minus the number of negative differences for all the differences considered. For large samples, the test is conducted using a normal distribution [46] with mean and variance as follows:

$$E[S] = 0 \quad (5)$$

$$\text{Var}(S) = \frac{N(N-1)(2N+5) - \sum_{k=1}^n t_k(t_k-1)(2t_k+5)}{18} \quad (6)$$

where, n is the number of tied (zero difference between compared values) groups and t_k is the number of data points in the k^{th} tied group. The standard normal deviate (Z -statistics) is then computed as :

$$Z = \begin{cases} \frac{S-1}{\sqrt{\text{Var}(S)}} & \text{if } S > 0 \\ 0 & \text{if } S = 0 \\ \frac{S+1}{\sqrt{\text{Var}(S)}} & \text{if } S < 0. \end{cases} \quad (7)$$

If the computed value of $|Z| > Z_{\alpha/2}$, the null hypothesis H_0 is rejected at the α level of significance in a two-sided test. In this analysis, the null hypothesis was tested at 95% confidence level.

3. RESULTS AND DISCUSSION

The Mann–Kendall (MK) test has been used for identification and testing the statistical significance of trend at confidence interval of 95%, prior to which data series of all the variables was checked for the presence of auto-correlation. The Sen’s estimator of slope (SE) was then applied to estimate the magnitude of the trend over the study period.

3.1 Trends in Temperature

In order to analyze trend, the anomalies of maximum temperature (T_{max}), minimum temperature (T_{min}), average temperature (T_{avg}) were computed firstly; and then the MK test was applied for the data of each station at both seasonal and annual scale. The results of the test for the increasing or decreasing trend along with their magnitude which was estimated using Sen’s slope estimator (SE) are given in Tables 2 to 4.

The results as presented in Table 2, shows an increasing trend in average temperature for all the four stations at annual as well as seasonal scale. The trend, however, was significant for annual and winter season scale. The magnitude of the annual trend for average temperature (T_{avg}) ranges from 0.020 °C/year (for Indore) to 0.024 °C/year (for Mahidpur). Similarly, magnitude of trend in the winter seasons was varies from 0.031 °C/year to 0.033 °C/year. In the post monsoon season the trend magnitude of average temperature was also significant except Alot station. Irrespective of scales (i.e. annual and seasonal), the average temperature for all the four station viz. Alot, Mahidpur, Indore and Dewas showed an increasing trend. For annual and winter temperature the rising trend is significant for all the four stations.

As seen from Table 3, the trend for maximum temperature (T_{max}) is also similar to that for average temperature (T_{avg}); however, the magnitude of annual trend for maximum temperature (T_{max}) ranges from 0.020 °C/year (for Indore) to 0.024 °C/year (for Mahidpur). Similarly, magnitude of trend in the winter seasons was varies from 0.033°C/year to 0.036

°C/year. Statistically significant increasing trend is found in post-monsoon except Alot station. The Mahidpur and Dewas station also showed the highest slope values for Tmax (0.041 °C/year) for post-monsoon season.

The trend for minimum temperature was also found to observe as increasing for all the four station for all time scales (Table 4). The trend however was significant for only annual and winter season scale. As seen from Table 4, the magnitude of annual as well as seasonal minimum temperature trend was relatively low as compared to maximum and average temperature. On average Mahidpur was the station who has observed highest rise in maximum/ minimum/ average temperature annually as well as seasonally.

3.2 Trends in Rainfall

The general characteristics of seasonal and annual rainfall for all the four stations in the Kshipra river basin are given in Table 5. The coefficient of variation (CV) of the annual rainfall varies from 20.2% (at Indore) to 27.4% (at Alot). It is evident from the Table 5 that all the station receives maximum rainfall during the monsoon season and least rainfall during the winter season. The contribution of monsoon rainfall varies from 91% (Indore) to 95% (Alot), while contribution of post-monsoon season almost 5%.

Similar to temperature, the anomalies of rainfall and their trends were determined for all the stations considered in the study. The magnitude of the seasonal and annual trend in the time series as determined using the Sen's estimator is given in Table 6. The annual rainfall indicates decreasing trend at two station namely Mahidpur, Dewas; and increasing trends at other two stations with maximum increase (3.12 mm/year) at Indore. The increasing trend at Alot is of the order of 0.19 mm/year. Seasonal analysis of rainfall trends shows that all stations during post-monsoon and winter season experienced decreasing trend whereas pre-monsoon season indicated increasing trend except Dewas station.

3.3 Homogeneity Test Analysis and Percentage Change of Rainfall and Temperature over a Period

Homogeneity test has been performed using long term data to find out the shift year, when a considerable change in the data series was observed. The shift year (break point) in the data series is calculated by Mann-Whitney-Pettit (MWP) test [47,48,49] and results are given in Table 7. As seen from this table, overall the year 2000 and 1993 can be considered as the break point in case of rainfall and temperature series, respectively. The graphical representation of shift in the annual maximum and average temperature is given in Figs. 2 and 3, respectively.

Table 2. Annual and Seasonal trend of average temperature (T_{avg})

Season	Station	Z _{MK}	S	Sen slope(SE)	Serial correlation	Trend
Annual	Alot	3.27	212	0.023	0.255	Rising
	Mahidpur	3.39	220	0.024	0.278	Rising
	Indore	3.21	208	0.020	0.269	Rising
	Dewas	3.55	230	0.022	0.253	Rising
Pre-monsoon (Mar-May)	Alot	1.01	66	0.015	0.133	Rising
	Mahidpur	1.13	74	0.016	0.164	Rising
	Indore	1.07	70	0.014	0.134	Rising
	Dewas	1.22	80	0.013	0.074	Rising
Monsoon (June- Sept.)	Alot	1.01	66	0.013	-0.039	Rising
	Mahidpur	1.07	70	0.013	-0.029	Rising
	Indore	0.70	46	0.009	-0.051	Rising
	Dewas	0.88	58	0.012	-0.056	Rising
Post-Monsoon (Oct-Nov.)	Alot	1.78	116	0.039	0.252	Rising
	Mahidpur	2.18	142	0.042	0.264	Rising
	Indore	2.22	144	0.040	0.251	Rising
	Dewas	2.43	158	0.042	0.265	Rising
Winter (Dec- Feb)	Alot	2.62	170	0.032	0.149	Rising
	Mahidpur	3.02	196	0.033	0.147	Rising
	Indore	3.08	200	0.031	0.158	Rising
	Dewas	3.08	200	0.033	0.169	Rising

(Bold shows the statistically significant trend at 5 % level)

Table 3. Annual and seasonal trend of maximum temperature (T_{max})

Season	Station	Z _{MK}	S	Sen slope(SE)	Serial correlation	Trend
Annual	Alot	3.15	204	0.022	0.248	Rising
	Mahidpur	3.36	218	0.024	0.266	Rising
	Indore	3.27	212	0.020	0.247	Rising
	Dewas	3.49	226	0.021	0.231	Rising
Pre-monsoon (Mar-May)	Alot	0.88	58	0.013	0.134	Rising
	Mahidpur	1.01	66	0.013	0.166	Rising
	Indore	0.85	56	0.012	0.150	Rising
	Dewas	1.01	66	0.011	0.099	Rising
Monsoon (June- Sept.)	Alot	0.95	62	0.014	-0.042	Rising
	Mahidpur	1.01	66	0.013	-0.030	Rising
	Indore	0.67	44	0.009	-0.054	Rising
	Dewas	0.95	62	0.010	-0.052	Rising
Post- Monsoon (Oct-Nov.)	Alot	1.87	122	0.039	0.231	Rising
	Mahidpur	2.22	144	0.041	0.236	Rising
	Indore	2.15	140	0.039	0.214	Rising
	Dewas	2.37	154	0.041	0.224	Rising
Winter (Dec- Feb)	Alot	2.62	170	0.032	0.167	Rising
	Mahidpur	3.11	202	0.036	0.170	Rising
	Indore	3.02	196	0.034	0.178	Rising
	Dewas	3.27	212	0.036	0.181	Rising

Table 4. Annual and Seasonal trend of minimum temperature (T_{min})

Season	Station	Z _{MK}	S	Sen slope(SE)	Serial correlation	Trend
Annual	Alot	2.43	158	0.017	0.077	Rising
	Mahidpur	3.42	222	0.023	0.106	Rising
	Indore	2.96	192	0.020	0.083	Rising
	Dewas	3.21	208	0.021	0.080	Rising
Pre-monsoon (Mar-May)	Alot	0.29	20	0.004	0.008	Rising
	Mahidpur	1.66	108	0.023	0.131	Rising
	Indore	1.44	94	0.019	0.083	Rising
	Dewas	1.5	98	0.017	0.035	Rising
Monsoon (June- Sept.)	Alot	1.38	90	0.015	-0.167	Rising
	Mahidpur	1.72	112	0.022	-0.075	Rising
	Indore	1.44	94	0.016	-0.115	Rising
	Dewas	1.6	104	0.019	-0.113	Rising
Post- Monsoon (Oct-Nov.)	Alot	1.87	122	0.033	0.038	Rising
	Mahidpur	1.81	118	0.033	0.019	Rising
	Indore	1.53	100	0.030	-0.014	Rising
	Dewas	1.78	116	0.035	-0.005	Rising
Winter (Dec- Feb)	Alot	2.06	134	0.029	0.041	Rising
	Mahidpur	2.34	152	0.034	0.021	Rising
	Indore	2.37	154	0.029	0.062	Rising
	Dewas	2.43	158	0.031	0.058	Rising

Table 5. Distribution of rainfall at different stations in Kshipra basin (1980-2012)



Station	Annual		Pre-monsoon		Monsoon		Post-monsoon		Winter	
	Mean	CV	Mean	% of annual	Mean	% of annual	Mean	% of annual	Mean	%of annual
Alot	882.1	27.4	8.3	0.9	837.6	95.0	31.2	3.5	5.0	0.6
Mahidpur	890.3	27.0	11.5	1.3	831.1	93.4	39.0	4.4	8.6	1.0
Indore	792.1	20.2	13.9	1.8	724.9	91.5	44.9	5.7	8.4	1.1
Dewas	868.6	22.2	11.6	1.3	806.2	92.8	39.0	4.5	11.8	1.4

Table 6. Seasonal and annual trends in rainfall for different stations in Kshipra basin

Season	Station	ZMK	S	Sen slope (SE)	Serial correlation	Trend
Annual	Alot	0.08	6	0.19	0.13	Rising
	Mahidpur	-0.26	-18	-1.51	-0.02	Falling
	Indore	0.98	64	3.12	0.14	Rising
	Dewas	-0.57	-38	-1.79	-0.13	Falling
Pre-monsoon (Mar - May)	Alot	0.73	48	0.13	0.09	Rising
	Mahidpur	2.31	150	0.36	0.26	Rising
	Indore	0.77	51	0.18	-0.09	Rising
	Dewas	-1.17	-73	-0.28	0.43	Falling
Monsoon (Jun - Sept)	Alot	0.20	14	0.77	0.16	Rising
	Mahidpur	-0.20	-14	-0.54	0.04	Falling
	Indore	1.29	84	4.14	0.18	Rising
	Dewas	-0.20	-14	-0.38	-0.10	Falling
Post-monsoon (Oct - Nov)	Alot	-1.49	-97	-0.53	0.11	Falling
	Mahidpur	-1.09	-71	-0.43	0.09	Falling
	Indore	-0.81	-53	-0.48	0.09	Falling
	Dewas	-2.23	-145	-1.18	0.09	Falling
Winter (Dec - Feb)	Alot	-1.67	-109	-0.10	0.00	Falling
	Mahidpur	-1.15	-75	-0.09	0.00	Falling
	Indore	-1.98	-129	-0.14	-0.06	Falling
	Dewas	-4.26	-274	-0.64	0.13	Falling

Table 7. Value of homogeneity test (Pettitt's test)

Station	Rainfall		Tmax		Tmin		Tavg	
	Pettitt's test		Pettitt's test		Pettitt's test		Pettitt's test	
	Shift year	P value						
Dewas	2000	0.401	1991	0.002	1993	0.005	1991	0.001
Indore	2006	0.177	1993	0.003	1993	0.006	1993	0.003
Alot	1999	0.895	1991	0.003	1993	0.059	1991	0.002
Mahidpur	2000	0.580	1993	0.002	1993	0.003	1993	0.002






Fig. 2. Shift change for average annual maximum temperature




Fig. 3. Shift change for annual average temperature




Fig. 4. Percentage change in seasonal average temperature over the period 1993-2012




Fig. 5. Percentage change in seasonal maximum temperature over the period 1993-2012




Fig. 6. Percentages change of seasonal minimum temperature for the period of 1993-2012




Fig. 7. Percentage change of seasonal rainfall for the period of 2000-2012




Fig. 8. Percentage change in rainfall and temperature on annual basis

3.4 Spatial Analysis of Rainfall and Temperature Data

The spatial interpolation technique [50,51] was employed to determine the spatial pattern of rainfall and temperature using ArcGIS 10.1. Here, the distribution of percentage change in annual and seasonal temperature as well as rainfall series was detected by considering shift year as base (or reference) year. Further, the distribution of these percentage change was interpolated by using the Inverse Distance Weighting (IDW) method to show their spatial distributions and results are given in Figs. 4 to 8. As seen from Fig. 4, the highest seasonal average temperature change was observed in post monsoon season. In contrast to it, the lowest change was seen to observe in monsoon season. Among all four stations, Mahidpur station was found to experience highest change in temperature in all the four seasons. Fig. 5 found to show the similar type of results as do the Fig. 4. In case of seasonal minimum temperature change, the highest change was seen to observe in winter season followed by post monsoon, monsoon and pre-monsoon (Fig. 6). The Fig. 7 shows the seasonal distribution of rainfall change where the highest change was seen in winter season followed by pre-monsoon, post-monsoon and monsoon. Dewas station was found to observed highest negative change in winter season.

On annual basis, rainfall was decreased at Alot, Mahidpur and Dewas stations, whereas it was increased at Indore station. The annual temperature change pattern was also same for all the four stations. On average, annual minimum, maximum and average temperature was found to increase by 1.5 % over the period 1993 to 2012.

4. CONCLUSIONS

The following conclusions can be drawn from the trend analysis of 32 years series of rainfall and temperature (maximum, minimum and mean) data carried out for annual as well as seasonal scale for Kshipra river basin, Madhya Pradesh, India.

1. An increase in annual and seasonal maximum, minimum and average temperature was observed at the all four stations.
2. The rise in trends of annual and winter temperature were significant for all the

four stations. Among all four stations, Mahidpur station was found to experience highest change in temperature in all the four seasons.

3. The magnitude of rise in annual minimum and minimum temperatures was varies from 0.017°C/year to 0.023°C and 0.020°C/year to 0.024°C/year, respectively.
4. On annual scale, a non-significant increasing trend in rainfall was observed at stations, Indore and Alot, whereas decreasing trend was observed at station, Mahidpur and Dewas. Moreover, the magnitude of the trend varies from -1.51 mm/year in Mahidpur to 3.12 mm/year in Indore.
5. The 32 years (1980-2012) data series of rainfall and temperature was analysed by means of Pettit's test, and the years 2000 and 1993 were obtained as the break point/change year for rainfall and temperature series, respectively.
6. On average, annual minimum, maximum and average temperature was found to increase by 1.5 % over the period 1993 to 2012.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Chandniha SK, Meshram SG, Adamowski JF, Meshram C. Trend analysis of precipitation in Jharkhand State, India: Investigating precipitation variability in Jharkhand State. *Theor Appl Climatol*. 2017;130:261-74.
2. Sonali P, Kumar DN. Review of trend detection methods and their application to detect temperature changes in India. *J Hydrol*. 2013;476:212-27.
3. Vinnikov KY, Groisman PY, Luginina KM. Empirical data on contemporary global climate changes (temperature and precipitation). *J Climate*. 1990;3(6): 662-77.
4. Thapliyal V, Kulshrestha SM. Climate changes and trends over India. *Mausam*. 1991;42(4):333-38.
5. Jones PD, Raper SCB, Bradley RS, Diaz HF, Kelly PM, Wigley TML. Northern

- Hemisphere surface air temperature variations: 1851-1984. *J Climate Appl Meteorol.* 1986;25(2):161-79.
6. Folland CK, Parker DE. Observed variations of sea surface temperature. In: *Climate-ocean interaction*. Springer Netherlands; 1990;21-52.
 7. IPCC.Summary for Policymakers [Pörtner HO, Roberts DC, Poloczanska ES, Mintenbeck K, Tignor M, Alegría A, Craig M, Langsdorf S, Löschke S, Möller V, Okem A, (eds.)]. In: *Climate Change 2022: Impacts, adaptation, and vulnerability. contribution of working group II to the sixth assessment report of the intergovernmental panel on climate change* [Pörtner HO, Roberts DC, Tignor M, Poloczanska ES, Mintenbeck K, Alegría A, Craig M, Langsdorf S, Löschke S, Möller V, Okem A, Rama B (eds.)]. Cambridge university press, cambridge, UK and New York, NY, USA. 2022;3-33.
DOI: 10.1017/9781009325844.001
 8. Beniston M, Diaz HF, Bradley RS. Climatic change at high elevation sites: An overview. *Clim Change.* 1997;36(3-4):233-51.
 9. Wibig J, Glowicki B. Trends of minimum and maximum temperature in Poland. *Clim Res.* 2002;20(2):123-33.
 10. Beniston M. Climatic change in mountain regions: A review of possible impacts. In: *Climate variability and change in high elevation regions: Past, present & future*. Springer Netherlands. 2003;5-31.
 11. Diaz HF, Grosjean M, Graumlich L. Climate variability and change in high elevation regions: past, present and future. Springer Netherlands. 2003;1-4.
 12. Rebetez M, Dobbertin M. Climate change may already threaten Scots pine stands in the Swiss Alps. *Theor Appl Climatol.* 2004;79(1-2):1-9.
 13. Vuille M, Bradley RS, Werner M, Healy R, Keimig F. Modeling $\delta^{18}O$ in precipitation over the tropical Americas: 1. Interannual variability and climatic controls. *J Geophys Res Atmos.* 2003;108(D6).
 14. Bhutiyani MR, Kale VS, Pawar NJ. Long-term trends in maximum, minimum and mean annual air temperatures across the Northwestern Himalaya during the twentieth century. *Clim Change.* 2007;85(1-2):159-77.
 15. Dimri AP, Ganju A. Wintertime seasonal scale simulation over Western Himalaya using RegCM3. *Pure Appl Geophys.* 2007;164(8-9):1733-46.
 16. Dash SK, Jenamani RK, Kalsi SR, Panda SK. Some evidence of climate change in twentieth-century India. *Clim Change.* 2007;85(3-4):299-321.
Duhan D, Pandey A. Statistical analysis of long term spatial and temporal trends of precipitation during 1901–2002 at Madhya Pradesh, India. *Atmos Res.* 2013;122:136-49.
 17. Nath S, Mathew A, Khandelwal S, Shekar PR. Rainfall and temperature dynamics in four Indian states: A comprehensive spatial and temporal trend analysis. *HydroRes.* 2023;6:247-54.
 18. Wu Y, Fang H, Huang L, Ouyang W. Changing runoff due to temperature and precipitation variations in the dammed Jinsha River. *J Hydrol.* 2020;582: 124500.
 19. Ekwueme BN, Agunwamba JC. Modeling the influence of meteorological variables on runoff in a tropical watershed. *Civil Eng J.* 2020;6(12):2344-51.
 20. Mondal SK, Tao H, Huang J, Wang Y, Su B, Zhai J, et al. Projected changes in temperature, precipitation and potential evapotranspiration across Indus River Basin at 1.5–3.0°C warming levels using CMIP6-GCMs. *Sci Total Environ.* 2021;789:147867.
 21. Li C, Hao J, Zhang G, Fang H, Wang Y, Lu H. Runoff variations affected by climate change and human activities in Yarlung Zangbo River, southeastern Tibetan Plateau. *Catena.* 2023;230: 107184.
 22. Hirsch RM, Slack JR, Smith RA. Techniques of trend analysis for monthly water quality data. *Water Resour Res.* 1982;18(1):107-21.
 23. Yu YS, Zou S, Whittemore D. Non-parametric trend analysis of water quality data of rivers in Kansas. *J Hydrol.* 1993;150(1):61-80.
 24. Kothyari UC, Singh VP. Rainfall and temperature trends in India. *Hydrol Process.* 1996;10(3):357-72.
 25. Fu G, Chen S, Liu C, Shepard D. Hydro-climatic trends of the Yellow river basin for

- the last 50 years. *Clim Change*. 2004;65(1-2):149-78.
26. Tebakari T, Yoshitani J, Suvanpimol C. Time-space trend analysis in pan evaporation over Kingdom of Thailand. *J Hydrol Eng*. 2005;10(3):205-15.
 27. Jhajharia D, Shrivastava SK, Sarkar D, Sarkar S. Temporal characteristics of pan evaporation trends under the humid conditions of northeast India. *Agric For Meteorol*. 2009;149(5):763-70.
 28. Kumar R, Singh RD, Sharma KD. Water resources of India. *Curr Sci*. 2005;89(5):794-811.
 29. Chattopadhyay S, Jhajharia D, Chattopadhyay G. Univariate modelling of monthly maximum temperature time series over northeast India: Neural network versus Yule-Walker equation based approach. *Meteorol Appl*. 2011;18(1):70-82.
Available:<http://doi.org/10.1002/met.211>
 30. Dinpashoh Y, Jhajharia D, Fakheri-Fard A, Singh VP, Kahya E. Trends in reference crop evapotranspiration over Iran. *J Hydrol*. 2011;399(3):422-33.
 31. Jain SK, Kumar V. Trend analysis of rainfall and temperature data for India. *Curr Sci*. 2012;102(1):37-49.
 32. Jain SK, Tyagi J, Singh V. Simulation of runoff and sediment yield for a Himalayan watershed using SWAT model. *J Water Resour Prot*. 2010;2(3):267.
 33. Singh A, Singh S, Nema AK, Singh G, Gangwar A. Rainfall-runoff modeling using MIKE 11 NAM model for Vinayakpur intercepted catchment, Chhattisgarh. *Indian J Dryland Agric Res Dev*. 2014;29(2):1-4.
 34. Meena PK, Khare D, Shukla R, Mishra PK. Long term trend analysis of mega cities in Northern India using Rainfall Data. *Indian J Sci Technol*. 2015;8(3):247.
 35. Dubey V, Panigrahi S, Vidyarthi VK. Statistical trend analysis of major climatic factors over Chhattisgarh State, India. *Earth Syst Environ*. 2023;7(3):629-48.
 36. Sen PK. Estimates of the regression coefficient based on Kendall's tau. *J Am Stat Assoc*. 1968;63(324):1379-89.
 37. Lettenmaier DP, Wood EF, Wallis JR. Hydro-climatological trends in the continental United States, 1948-88. *J Climate*. 1994;7(4):586-607.
 38. Yue S, Hashino M. Long term trends of annual and monthly precipitation in Japan. *JAWRA J Am Water Resour Assoc*. 2003;39(3):587-96.
 39. Partal T, Kahya E. Trend analysis in turkish precipitation data. *Hydrol Process*. 2006;20(9):2011-26.
 40. Yu YS, Zou S, Whittemore D. Non-parametric trend analysis of water quality data of rivers in Kansas. *J Hydrol*. 1993;150(1):61-80.
 41. Burn DH, Cunderlik JM, Pietroniro A. Hydrological trends and variability in the Liard River basin / Tendances hydrologiques et variabilité dans le bassin de la rivière Liard. *Hydrol Sci J*. 2004;49(1):53-67.
Available:<http://doi.org/10.1623/hysj.49.1.53.53994>
 42. Neeti N, Eastman JR. A contextual Mann-Kendall approach for the assessment of trend significance in image time series. *Trans GIS*. 2011;15(5):599-611.
 43. Sayemuzzaman M, Jha MK. Seasonal and annual precipitation time series trend analysis in North Carolina, United States. *Atmos Res*. 2014;137:183-94.
 44. Kisi O, Ay M. Comparison of Mann-Kendall and innovative trend method for water quality parameters of the Kizilirmak River, Turkey. *J Hydrol*. 2014;513:362-75.
 45. Helsel DR, Hirsch RM. *Statistical methods in water resources*. Vol. 49. Elsevier; 1992.
 46. Hirsch RM, Slack JR, Smith RA. Techniques of trend analysis for monthly water quality data. *Water Resour Res*. 1982;18(1):107-21.
 47. Ngongondo CS. An analysis of long-term rainfall variability, trends and groundwater availability in the Mulunguzi river catchment area, Zomba mountain, Southern Malawi. *Quat Int*. 2006;148(1):45-50.
 48. Goyal MK. Statistical analysis of long term trends of rainfall during 1901-2002 at Assam, India. *Water Resour Manag*. 2014;28(6):1501-15.
 49. Bližňák V, Valente MA, Bethke J. Homogenization of time series from Portugal and its former colonies for the period from the late 19th to the early 21st century. *Int J Climatol*. 2015;35(9):2400-18.

50. Singh VP, Chowdhury PK. Comparing some methods of estimating mean areal rainfall. JAWRA J Am Water Resour Assoc. 1986;22(2):275-83.
51. Lebel T, Bastin G, Obled C, Creutin JD. On the accuracy of areal rainfall estimation: A case study. Water Resour Res. 1987;23(11):2123-34.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/118265>