



Medicinal Properties of Terminalia Arjuna: A Review

Vaishnavi Verma ^{a*} and Sangita Jogdand ^b

^a *Jawaharlal Nehru Medical College, Datta Meghe Institute of Medical Sciences, Sawangi (M), Wardha, Maharashtra, India.*

^b *Department of Pharmacology, Jawaharlal Nehru Medical College, Datta Meghe Institute of Medical Sciences, Sawangi (M), Wardha, Maharashtra, India.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i62A35893

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/80456>

Review Article

Received 28 October 2021
Accepted 26 December 2021
Published 28 December 2021

ABSTRACT

Terminalia arjuna, also commonly referred to as *T. arjuna*, is a deciduous tree that belongs to the family Combretaceae. It can be found in many regions of India. *T. arjuna* is a 60- to 80-foot-tall tree found alongside rivers and streams all over the Indo-sub-Himalayan areas of Delhi, Uttar Pradesh, Chota Nagpur, the southern part of Bihar, Madhya Pradesh and Deccan regions. It has been used to cure several ailments for as far back as the ancient times of India. It is most prevalently consumed to cure and manage several cardiac and vascular diseases, including those like CADs, Angina Pectoris, CHF/Hypertension, and Dyslipidaemia. Its extracts are used to improve cardiac muscles and thus effectively improve heart pumping, heart rate, and blood pressure. The many parts of the tree consist of several phytochemicals, including tannins, flavonoids, glycosides, and triterpenoids like Arjunolic acid, which contribute to its anti-oxidant anti-inflammatory antimicrobial, anticarcinogenic and antimutagenic properties. As of today, there have not been any reports of any harmful side effects regarding its administration. While there are various studies that support its use for a problem of diseases, further research is still required to understand its exact mechanisms. There is also a need for further research on *T. arjuna* regarding its drug interactions, its specific molecular mechanism of action, and the toxicology involved.

MBBS Student,

† Professor,

*Corresponding author: E-mail: vaishnavi26700@gmail.com;

Keywords: Chemical constituents; medicinal properties; cardiovascular; CAD; angina pectoris; CHF; tannins; flavonoids; triterpenoids arjunolic acid.

1. INTRODUCTION

Plants with therapeutic characteristics have long been utilized to heal ailments for thousands of years. [1]. WHO statistics estimate that approximately 80% of the world's population, including 60% of India's rural population, depend on these therapeutic agents [2]. The demand for herbal medicines has only increased in recent years. Due to their easy accessibility, efficiency, and rare side effects. These medicinal plants contain certain bioactive substances such as alkaloids, tannins, carbohydrates, steroids, terpenoids, phenols, and flavonoids, ensuring certain physiological effects on the body [3].

A variety of medicinal plants have been employed in modern-day healthcare, including:

Table 1. Drugs derived from plants [4]

Natural Plant Source	Name of the Drug
Foxglove	Digitalis
Willow Bark	Salicylates
Cinchona	Quinine
Contaminated Rye	Ergotamine

There are a variety of medicinal plants in India which have been extensively employed in Ayurvedic practices. *T. arjuna* is one of these plants, and it has proven to be one of the most commonly acknowledged herbal medicines for the treatment of a range of disorders [1].

1.1 *T. arjuna*: Overview

T. arjuna of the family Combretaceae. Long used as a cardioprotective agent, it was first introduced by Vagabhatta, who advocated its stem bark powder's use for heart diseases and has since been written in various ancient texts like the Sushruta Samhita, Charaka Samhita, and Ashtang Hridayam. [5].

T. arjuna is a 60- to 80-foot-tall tree found alongside rivers and streams all over the Indo-sub-Himalayan areas of Delhi, Uttar Pradesh, Chota Nagpur, the southern part of Bihar, Madhya Pradesh, and Deccan regions. Aside from being found in various regions of India, it has also been seen in many other countries, including Sri Lanka, Burma as well as Mauritius [6,7]. Although this plant grows on all types of

soil, it has shown a preference for red lateric, fertile loam and humid soil.

The tree's bark has a smooth outer surface and an inner striated pinkish surface [8]. During April and May, the bark of this tree sheds away [9].



Fig. 1. The bark of the tree *Terminalia arjuna*

1.2 Chemical Components of *Terminalia arjuna*

The learoots roots, fruits, ste, as well as ieedes of *T.arjuna*, have been used in medical practice due to their different phytoconstituents.

2. TERPENOIDS, URSANE TRITERPENOIDS AND GLYCOSIDES

Triterpenoids are structurally diverse organic compounds which include various varieties, due to modifications in its basic backbone, including ursolic and oleanolic acid [39].

The table above lists a number of terpenoids, ursane triterpenoids, and glycosides isolated from a variety of areas of *T. arjuna*.

Each type has its own pharmacodynamic effect on the body. Ali et al [24] discovered Terminoside A, an oleanane-type triterpane, from *T. arjuna* stem bark in a research. The Terminoside A thus extracted exhibited characteristics that prevented the synthesis of nitric oxide. In macrophages stimulated by lipopolysaccharides, it also decreased the quantity of inducible nitric oxide synthase (iNOS or simply iNOS [20,21].

Table 2. Phytochemical components of various sections of *T. arjuna* [1]

Parts of the tree that were analyzed	Chemical components which are considered important
Stem Bark	<p>Ursane triterpenoids 2α,3β-dihydroyurs-12,18-oic acid 28-O-β-D-glucopyranosyl ester [10] Kajiichigoside F1 2α,3β,23-trihydroxyurs-12,18-dien-28-oic acid 28-O-β-glucopyranosyl ester Qudranoside VIII 2α,3β,23-trihydroxyurs-23-trihydroxyurs-12,19-dien-28-oic acid 28-O-β-D-glucopyranosyl ester</p> <p>Triterpenoids Arjunin [11] Arjunic acid Arjungenin [12-14] Terminic acid [15] Arjunolic acid [13,14] Terminoltin [16]</p> <p>Flavonoids and Phenolics Arjunone [17] Luteolin [18] Baicalein [19] Ethyl gallate Kempferol Gallic acid Pelargonidin Oligomeric proanthocyanidins Quercetin Gallic acid, ellagic acid and its derivatives such as 3-O-methyl-ellagic acid 4-O-β-D-xylopyranoside, 3-O-methyl ellagic acid 3-O-rhamnoside (+)-catechin, (+)-gallocatechin and (-)-epigallocatechin [20] 3-O-methyl ellagic acid 4'-O-α-L-rhamnophranoside (-)-epicatechin [10]</p> <p>Glycosides Arjunetin [11,21,13,14] Arjunolone [17] Arjunoside I, II [12,22; Arjunaphthanoloside [23,24] Arjunolitin [25] Arjunasides A-E, Arjunglucoside IV and V [25,26] Terminarjunoside I and II [27] Olean-3β, 22β-diol-12-en-28 β-D-glucopyranosie-oic acid [28] Terminoside A [29] Termionic acid</p> <p>Trace elements along with Minerals Magnesium, Calcium, aluminium, silica, zinc, copper [30]</p> <p>Tannins Pyrocatechols [31] Castalagin [32] Punicallin [33] Casuariin Casuarinin Punicalagin Terflavin C</p>

Parts of the tree that were analyzed	Chemical components which are considered important
Roots	Terchebulin Other compounds β-Sitosterol [15] Glycosides Arjunetosie (3-O-β-D-glucopyranosyl-2α, 3β, 19α-trihydroxyolean-12-en-28-oic acid 28-O-β-D-glucopyranoside) [34] Triterpenoids Arjunoside I-IV [35] Oleanolic acid Arjunolic acid [15] 2α,19α-Dihydroxy-3Oxo-Olean-12-En28-Olic acid 28-O-β-D-glucopyranoside [36] Terminic acid Arjunic acid [13,14]
Seeds and Leaves	Glycosides and Flavonoids Luteolin, 14,16-dianhydrogitoxigenin 3-β-D-xylopyranosyl-(1 > 2)-O-β-D-galactopyranoside [18,37]
Fruits	Flavonoids and Triterpenoids Hentriacontane, Arjunic acid, Ellagic acid, Arjunone, Eridelin, Methyl oleolate, Gallic acid, Cerasidin, Myristyl oleate, β-Sitisterol, Arachidic stearate [38]

3. FLAVONOIDS ALONG WITH PHENOLICS

From a medicinal point of view, *T. arjuna*'s bark is perhaps the most significant portion of *T. arjuna*'s bark is regarded the most important part of the plant. The bark contains a variety of flavonoids such as flavones, arjunolones, kempferol, baicalein, pelargonidin and quercetin. Because of an inverse link between high dietary flavonoid consumption and the development of ischemic heart diseases (CADs), these flavonoids are particularly useful for treating cardiovascular disorders [1].

Luteolin, a molecule isolated from the butanolic fraction, exhibits antimutagenic properties [1]. It also has a very efficient antibacterial property as it inhibits gram negative pathogenic growth with the minimum inhibitory concentration of 12.5µg/disc.

Other actions of these bioflavonoids include inhibition of oxidation of LDL molecules, activation of endothelium and aggregation of platelets [1,40-43].

The phenolic content contributes to a free radical scavenging action which makes *T. arjuna* a strong agent against proliferation and oxidation [1,44].

Tannins: Tannins are polyphenols that are water soluble and may be found in a range of plant components. Tannins possess a variety of properties. One such property is that it is an anticarcinogen along with tea polyphenols. It also has an anti-mutagenic property as well as an anti-oxidant property. These three properties are interrelated as oxygen-free radicals are produced by a variety of carcinogens and mutagens. which these tannins ultimately decrease for protecting cellular oxidative cellular damage.

Another important property of tannins is its antimicrobial activity. Studies have shown that Yeasts, fungi, bacteria, and viruses have all been found to be inhibited by tannins.

Tannins also aid in the clotting of blood, the reduction of blood pressure, the reduction of serum cholesterol levels, the production of liver necrosis, and the modulation of immune responses [45].

4. PHARMACOLOGICAL ACTIVITIES OF *Terminalia arjuna*

Even though each part of *T. arjuna* has its own pharmacological effects on the body due to their varying composition, the bark of the tree is regarded to the most clinically relevant.

The bark has been shown to have astringent, expectorant, demulcent, cardiogenic, anti-dysenteric, styptic and urinary astringent effects, as well as being beneficial in the management of cirrhosis, anaemia, leukorrhea, fractures, cardiomyopathy, diabetes and ulcers [8,46].

Chakradatta introduced an ulcer wash made from an infusion of the bark prepared using milk and perhaps even ghee/butter in Ancient India. The ashes of the bark was employed for the management of snakebites and scorpion stings [47].

It has been used in many forms throughout India for a variety of conditions. *T. arjuna*'s bark is boiled in water and breathed in which then alleviates headaches and eliminate worms in the teeth in the Kancheepuram District of Kerala. They also use its fruit's paste as a topical agent on wounds [48]. The bark powder is mixed with rice water by tribals in the Sundargarh District of Odisha to treat haematuria (there is blood in the urine [49].

A more detailed analysis of the same on the basis of clinical trials and experiments has been tabulated below.

Table 3. *Terminalia arjuna* – pharmacological activities [1]

Pharmacological activity	Chemical Constituents Responsible	Supporting Clinical Trial/Experiment	Observation in the concerned clinical trial/experiment
Antioxidant, anti-inflammatory and immunomodulatory	Arjunic acid, arjunetin and arjungenin	Varghese et al. [50]	The enzymes were shown to have strong non-competitive inhibitory and reversible action in both of <i>T. arjuna</i> 's aqueous and alcoholic extracts. The enzymes concerned are CYP3A4, CYP2D6 and CYP2C9 present in the human liver microsomes [1].
Antioxidant	Oleanane triterpenoids	Pawar and Bhutani [54]	The process of respiratory burst is modestly inhibited by Arjungenin. Its IC50 results to be 60 µg/ml [1]
Antioxidant activity	Butanolic fraction of <i>Terminalia arjuna</i> bark [52]	Singh et al [52]	The butanolic component of <i>T. arjuna</i> 's bark's alcoholic extract shows cardioprotective activity in a patient with Doxorubicin-induced cardiotoxicity.
Antioxidant and antimutagenic activity	Alcoholic extract of <i>Terminalia arjuna</i> stem bark (ALTA)	Viswanatha et al [53]	In the DPPH assay, liquid peroxidation assay and superoxide radical scavenging activity, the alcoholic extract of <i>T. arjuna</i> showed significant antioxidant activity with EC50 values of 2.491 ± 0.160, 71.000 ± 0.025, and 50.110 ± 0.150 respectively. In the micronucleus test, EC50 values of 2.410 ± 0.140, 40.500 ± 0.390, and 63.000 ± 0.360 in percentage of micronucleus in <i>T. arjuna</i> 's alcoholic extract (100 and 200 mg/kg p.o) resulted in significant reductions in both the normochromatic and polychromatic erythrocytes, as well as quite a decrease in the P/N ratio [1]
Potential to be antimutagenic and anticarcinogenic	<i>Terminalia arjuna</i> bark has substantial flavonoids and tannins.	Ahmad et al [54]	<i>T. arjuna</i> extracts were shown to be effective in reducing metaphase abnormalities. In vitro, the frequency of sister chromatid exchanges was decreased, but the replication index rose. Clastogeny was reduced in the

Pharmacological activity	Chemical Constituents Responsible	Supporting Clinical Trial/Experiment	Observation in the concerned clinical trial/experiment
			mutagen-treated positive control and aberrant cell frequencies were reduced in the in vivo trials.
Anti-oxidant and antimicrobial activity	<i>Terminalia arjuna</i> bark's Methanolic extract	Mandal et al. [55]	<i>T.arjuna's</i> methanolic extracts have potent antibacterial activity as well as scavenging of free radicals. It is a potent antibacterial agent against <i>K. pneumonia</i> and <i>E. coli</i> (gram-negative bacteria). These properties are because of the flavonoid compounds in <i>T. arjuna</i> [56]
Antimicrobial activity	<i>Terminalia arjuna</i> leaf extract (acetonic) and bark extract (aqueous [7]	Aneja et al. [57]	The aqueous extract resulted in being an efficient antimicrobial against <i>S. aureus</i> bacteria [58] However, the acetone extract of the leaf extract of <i>T. arjuna</i> was seen to have the most potent antimicrobial agent against <i>S. aureus</i> [1]. The organic extracts were shown to be highly efficient against the proliferation of gram-negative bacteria, with the exception of <i>P. aeruginosa</i> .
Cardio-protective potential	<i>Terminalia arjuna</i> bark powder which was used for 12 weeks before ischemic-reperfusion injury [1]	Gauthaman et al [59].	Myocardial endogenous antioxidants were boosted following chronic oral treatment of <i>T. arjuna</i> bark in rabbits. It also induces HSP-72 (Inducible Heat Shock Protein 72). The prevents myocardial ischemic reperfusion injuries due to protection against oxidative stress.
Anticarcinogenic potential	<i>Terminalia arjuna's</i> ethanolic and aqueous extracts	Oberoi et al. [60]	<i>T. arjuna's</i> aqueous extracts enhances sarcoplasmic reticular function and thus induces cardiogenic action. Arrhythmias are less likely to arise as a result of this. As a result, <i>T. arjuna's</i> aqueous extract is seems as a safe cardiogenic that is good to heart health and may be used in conjunction with chronic health-care treatment programmes [1]
Free radical scavenging and DNA damage protection	<i>Terminalia arjuna</i> bark's ethanolic extract along with its fractions	Phani Kumar et al. [61]	<i>T. arjuna</i> bark's ethanolic extract (together with its components) protect against hydrogen peroxide-induced DNA damage [62] The ethyl acetate fraction has especially been effective in maximally inhibiting DPPH, ABTS, metal chelation, hydroxyl and nitric oxide radicals. <i>T. arjuna</i> extracts have also been demonstrated to ameliorate a variety of impairments related to free radical production and DNA damage.

Pharmacological activity	Chemical Constituents Responsible	Supporting Clinical Trial/Experiment	Observation in the concerned clinical trial/experiment
Gastro-productive effect	Methanolic extract of <i>Terminalia arjuna</i>	Devi et al. [63]	Two groups of ulcer-induced animals were studied. One group received Diclofenac Sodium (DIC) and <i>T. arjuna</i> , whereas the other received simply Diclofenac Sodium. In comparison to merely providing DIC, the DIC + <i>T. arjuna</i> treatment plan demonstrated a considerable reduction in the lesion index [64]. <i>T. arjuna</i> 's gastroprotective effect was validated by other histological research.

5. CARDIOVASCULAR ROLE OF *T. arjuna*

The bark stem of *T. arjuna* has inotropic, chronotropic and diuretic properties [7]. Experiments on animals revealed an augmentation in coronary blood flow, which increased the force of cardiac muscle contraction, resulting in a drop in blood pressure along with heart rate as well as bradycardia with accordance with the dose administered [65,66,67,7].

Research done on rats found that pretreatment with atropine reduced the hypotensive effect of *T. arjuna* with a fraction containing tannin-related chemicals isolated from the aqueous extract. Pretreatment of the rats with propranolol had no impact, suggesting that the hypotensive effect was related to cholinergic processes [31].

In myocardial infarction which is induced by isoprenaline, *T. arjuna* exhibited PGE₂ like activity in the heart by producing vasodilatation and hypotension [5]. *T. arjuna*'s bark extract reduced the oxidative stress which upsurged on induction by isoprenaline and reduced the amount of natural antioxidants in the body [5].

One of the triterpenoids found in *T. arjuna*, Arjunolic acid, prevents the decline of superoxide dismutase, glutathione peroxidase, catalase, alpha-tocopherol, ceruloplasmin, ascorbic acid, reduced glutathione (GSH), MPO (myeloperoxidase) and lipid peroxide levels, implying that Arjunolic acid's cardioprotection by Arjunolic acid is most likely due to protection against damage to heart via myocardial necrosis [68]. Another study found that arjunolic acid had cardioprotective properties through boosting the body's natural antioxidant defences [69].

Animal experiments showed that the *T. arjuna* bark, when administered in various forms, was capable of reducing total cholesterol (TC) and triglyceride (TG) levels [5,70,71,72,73].

When compared to the other fractions of *T. arjuna* bark, the ethanolic fraction has powerful antioxidant and hypolipidemic effects [74,75]. The down regulation of lipogenic enzymes, the enhances hepatic clearance of cholesterol and inhibition of HMG-CoA reductase are likely to be responsible for the hypolipidemic effect [76].

Angina Pectoris: A study was undertaken where the sample size of 30 patients suffering from stable angina were administered with 500 mg of *T. arjuna* bark extract three times a day. The bark's anti-ischemic activity was proven by a considerable reduction in the serum cholesterol levels, systolic blood pressure, plasma cortisol and the mean anginal frequency. There was also an improvement in the ECG changes [5,77].

Chf/ Hypertension: A study was undertaken where the sample size of 10 patients suffering from Congestive Heart Failure (CHF) were given 4g of *T. arjuna* bark powder twice a day for a month as part of a research. With considerable diuresis, there was enhancement seen with dyspnea, functional class, and general well-being. Both the systolic as well as the diastolic blood pressures dropped significantly [78].

6. TOXICITY AND SIDE EFFECTS OF *Terminalia arjuna*

Most traditional and herbal medicines like *T. arjuna* are known for producing the least amount of side effects, hence their popularity. No cases of *T. arjuna* toxicity have been documented [19].

T. arjuna is most widely used for the cure and control of coronary artery disorders (CAD), with an ideal dose of 1-2 g per day, and 500 mg of the bark extract three times per day for congestive heart failure. The side effects reported in this treatment are rather minor like headaches, mild gastritis and constipation. After more than 2 years of this drug administration, there were no signs of haematological, hepatic, metabolic and renal toxicity [77,79].

A study reported that there was a reduction in thyroid hormone concentration in euthyroid animals and an increase in hepatic LPO (Lipid Peroxidation) upon the administration of *T. arjuna*. Therefore, care must be taken when consuming this plant extract as it carries a risk of development of hypothyroidism and hepatotoxicity [80-85].

7. CONCLUSION

T. arjuna, a tree seen all around India, is being utilised for hundreds of years for curing a conundrum of ailments, but more importantly for cardiac health. Its active constituents include tannins, triterpenoids, flavonoids and certain minerals like calcium, magnesium, zinc and copper.

Its extracts are used for the improvement of cardiac muscles, effectively improving heart pumping, heart rate and blood pressure.

Terminalia arjuna can be administered in a variety of conditions such as Angina Pectoris, Congestive Heart Failure, Cardiomyopathy or Post Myocardial Infarction and Hyperlipidemia.

While there are various studies which support their application in clinical practice, such studies lack the standardisation of extract to be used, well conducted studies for long term effects and the bioavailability of the drug.

There is also a need for further research on *T. arjuna* regarding its drug interactions, its specific molecular mechanism of action as well as toxicology.

NOTE

The study highlights the efficacy of "herbal, ayurvedic, traditional" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in

the light of modern medical science and can be utilized partially if found suitable.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Amalraj A, Gopi S. Medicinal properties of *Terminalia arjuna* (Roxb.) Wight & Arn.: A review. *J Tradit Complement Med.* 2016;7(1):65–78.
2. World Health Organization. WHO Traditional medicine strategy Report. Document WHO/EDM/TRM; 2002.
3. Sharma J, Gairola S, Gaur RD, Painuli RM. The treatment of jaundice with medicinal plants in indigenous communities of the Sub-Himalayan region of Uttarakhand, India. *J Ethnopharmacol.* 2012;143(1):262–91.
4. Sen T, Samanta SK. Medicinal Plants, Human Health and Biodiversity: A Broad Review. In: Mukherjee J, editor. *Biotechnological Applications of Biodiversity* [Internet]. Berlin, Heidelberg: Springer Berlin Heidelberg; 2014 [cited 2021 Aug 25]:59–110. Available: http://link.springer.com/10.1007/10_2014_273
5. Dwivedi S, Chopra D. Revisiting *Terminalia arjuna* – An Ancient Cardiovascular Drug. *J Tradit Complement Med.* 2014;4(4):224–31.
6. Chopra RN, Chopra IC. *Indigenous drugs of India.* Academic publishers; 1994.
7. Dwivedi S. *Terminalia arjuna* Wight & Arn.—A useful drug for cardiovascular disorders. *J Ethnopharmacol.* 2007;114(2): 114–29.
8. Warrier PK. *Indian medicinal plants: a compendium of 500 species.* Orient Blackswan; 1993.
9. Dhingra V, Dhingra S, Singla A. *Forensic and pharmacognostic studies of the*

- Terminalia arjuna Bark. Egyptian Journal of Forensic Sciences. 2013;3(1):15-9.
10. Wang W, Ali Z, Shen Y, Li XC, Khan IA. Ursane triterpenoids from the bark of Terminalia arjuna. Fitoterapia. 2010;81(6):480-4.
 11. Row LR, Murty PS, Rao GS, Sastry CP, Rao KV. Chemical examination of Terminalia species. XII. Isolation & structure determination of arjunic acid, a new trihydroxytriterpene carboxylic acid from Terminalia arjuna bark. Indian journal of chemistry; 1970.
 12. Honda T, Murae T, Tsuyuki T, Takahashi T, Sawai M. Arjungenin, arjunglucoside I, and arjunglucoside II. A new triterpene and new triterpene glucosides from Terminalia arjuna. Bulletin of the Chemical Society of Japan. 1976;49(11):3213-8.
 13. Singh DV, Verma RK, Gupta MM, Kumar S. Quantitative determination of olean derivatives in Terminalia arjuna by high performance thin layer chromatography. Phytochem Anal. 2002;13(4):207-10.
 14. Singh DV, Verma RK, Singh SC, Gupta MM. RP-LC determination of olean derivatives in Terminalia arjuna. J Pharm Biomed Anal. 2002;28(3-4):447-52.
 15. Anjaneyulu AS, Prasad AR. Structure of terminic acid, a dihydroxytriterpene carboxylic acid from Terminalia arjuna. Phytochemistry. 1983;22(4):993-8.
 16. Singh B, Singh V, Pandey V, Rücker G. A New Triterpene Glycoside from Terminalia arjuna. Planta Med. 1995;61(06):576-7.
 17. Sharma PN, Shoeb A, Kapil RS, Popli SP. Arjunolone-a New flavone from stem bark of terminalia-arjuna. Indian Journal of Chemistry Section b-Organic chemistry Including Medicinal Chemistry. 1982;21(3): 263-4.
 18. Pettit GR, Hoard MS, Doubek DL, Schmidt JM, Pettit RK, Tackett LP, et al. Antineoplastic agents 338. The cancer cell growth inhibitory. Constituents of Terminalia arjuna (Combretaceae). J Ethnopharmacol. 1996;53(2):57-63.
 19. Terminalia arjuna. Altern Med Rev J Clin Ther. 1999;4(6):436-7.
 20. Jayaraman S, Saha A, Pawar V. Characterisation of polyphenols in Terminalia arjuna bark extract. Indian J Pharm Sci. 2012;74(4):339.
 21. Row LR, Murti PS, Rao GS, Sastry CS, Rao KV. Chemical examination of Terminalia species. XIII. Isolation and structure determination of arjunetin from Terminalia arjuna. Indian journal of chemistry; 1970.
 22. Honda T, Murae T, Tsuyuki T, Takahashi T. The structure of arjungenin. A new sapogenin from Terminalia arjuna. Chemical and Pharmaceutical Bulletin. 1976;24(1):178-80.
 23. Ali A, Kaur G, Hayat K, Ali M, Ather M. A novel naphthanol glycoside from Terminalia arjuna with antioxidant and nitric oxide inhibitory activities. Pharm. 2003;58(12):932-4.
 24. Ali A, Kaur G, Hamid H, Abdullah T, Ali M, Niwa M, et al. Terminoside A, a new triterpene glycoside from the bark of Terminalia arjuna inhibits nitric oxide production in murine macrophages. J Asian Nat Prod Res. 2003;5(2):137-42.
 25. Tripathi VK, Pandey VB, Udupa KN, Ru G. Arjunolitin, a triterpene glycoside from Terminalia arjuna. Phytochemistry. 1992;31(1):349-51.
 26. Wang W, Ali Z, Li X-C, Shen Y, Khan IA. Triterpenoids from two Terminalia species. Planta Med. 2010;76(15):1751-4.
 27. Sarwar Alam M, Kaur G, Ali A, Hamid H, Ali M, Athar M. Two new bioactive oleanane triterpene glycosides from Terminalia arjuna. Nat Prod Res. 2008; 22(14):1279-88.
 28. Patnaik T, Dey RK, Gouda P. Isolation of triterpenoid glycoside from bark of Terminalia arjuna using chromatographic technique and investigation of pharmacological behavior upon muscle tissues. E-Journal of Chemistry. 2007 4(4):474-9.
 29. Ahmad MU, Mullah KB, Norin T, Ulla JK. Terminic acid, a new trihydroxytriterpene carboxylic-acid from bark of terminalia-arjuna. Indian Journal of Chemistry Section B-Organic Chemistry Including Medicinal Chemistry. 1983;22(8): 738-40.
 30. Dwivedi S, Udupa N. Terminalia arjuna: pharmacognosy, phytochemistry, pharmacology and clinical use. A review. Fitoterapia. 1989;60(5):413-20.
 31. Takahashi S, Tanaka H, Hano Y, Ito K, Nomura T, Shigenobu K. Hypotensive effect in rats of hydrophilic extract from Terminalia arjuna containing tannin-related compounds. Phytotherapy Research: An International Journal Devoted to Medical and Scientific Research on Plants and Plant Products. 1997;11(6):424-7.
 32. Kuo PL, Hsu YL, Lin TC, Lin LT, Chang JK, Lin CC. Casuarinin from the bark of

- Terminalia arjuna induces apoptosis and cell cycle arrest in human breast adenocarcinoma MCF-7 cells. *Planta Med.* 2005;71(3):237–43.
33. Lin TC, Chien SC, Chen HF, Hsu FL. Tannins and related compounds from Combretaceae plants. *Chinese Pharmaceutical Journal.* 2000;52(1):1-26.
 34. Upadhyay RK, Pandey MB, Jha RN, Singh VP, Pandey VB. Triterpene glycoside from Terminalia arjuna. *J Asian Nat Prod Res.* 2001;3(3):207–12.
 35. Anjaneyulu AS, Prasad AR. Chemical examination of the roots of Terminalia arjuna—the structures of arjunoside III and arjunoside IV, two new triterpenoid glycosides. *Phytochemistry.* 1982;21(8):2057-60.
 36. Chouksey BK, Srivastava SK. Antifungal agent from Terminalia arjuna. *Indian J Chem. B.* 2001;40:354-6.
 37. Yadava RN, Rathore K. A new cardenolide from the seeds of Terminalia arjuna (W&A). *J Asian Nat Prod Res.* 2000;2(2):97–101.
 38. Rastogi RP, Mehrotra BN. CSIR; New Delhi: (Compendium of Indian Medicinal Plants). 1993;3.
 39. Petronelli A, Pannitteri G, Testa U. Triterpenoids as new promising anticancer drugs. *Anticancer Drugs.* 2009;20(10):880–92.
 40. Fuhrman B, Aviram M. Anti-atherogenicity of nutritional antioxidants. *IDrugs Investig Drugs J.* 2001;4(1):82–92.
 41. Carluccio MA, Siculella L, Ancora MA, Massaro M, Scoditti E, Storelli C, et al. Olive oil and red wine antioxidant polyphenols inhibit endothelial activation: antiatherogenic properties of Mediterranean diet phytochemicals. *Arterioscler Thromb Vasc Biol.* 2003 Apr 1;23(4):622–9.
 42. Ruf JC. Wine and polyphenols related to platelet aggregation and atherothrombosis. *Drugs Exp Clin Res.* 1999;25(2–3):125–31.
 43. Martikainen JA, Ottelin A-M, Kiviniemi V, Gylling H. Plant stanol esters are potentially cost-effective in the prevention of coronary heart disease in men: Bayesian modelling approach. *Eur J Cardiovasc Prev Rehabil Off J Eur Soc Cardiol Work Groups Epidemiol Prev Card Rehabil Exerc Physiol.* 2007;14(2):265–72.
 44. Bajpai M, Pande A, Tewari SK, Prakash D. Phenolic contents and antioxidant activity of some food and medicinal plants. *Int J Food Sci Nutr.* 2005;56(4):287–91.
 45. Chung KT, Wong TY, Wei CI, Huang YW, Lin Y. Tannins and human health: a review. *Crit Rev Food Sci Nutr.* 1998;38(6):421–64.
 46. Greco G, Turrini E, Tacchini M, Maresca I, Fimognari C. The Alcoholic Bark Extract of Terminalia Arjuna Exhibits Cytotoxic and Cytostatic Activity on Jurkat Leukemia Cells. *Venoms Toxins.* 2021;1(1):56–66.
 47. Jain S, Yadav PP, Gill V, Vasudeva N, Singla N. Terminalia arjuna a sacred medicinal plant: phytochemical and pharmacological profile. *Phytochemistry Reviews.* 2009;8(2):491-502.
 48. Muthu C, Ayyanar M, Raja N, Ignacimuthu S. Medicinal plants used by traditional healers in Kancheepuram district of Tamil Nadu, India. *J Ethnobiol Ethnomedicine.* 2006;2:43.
 49. Prusti AB, Behera KK. Ethnobotanical exploration of Malkangiri district of Orissa, India. *Ethnobotanical leaflets.* 2007; (1):14.
 50. Varghese A, Savai J, Pandita N, Gaud R. In vitro modulatory effects of Terminalia arjuna , arjunic acid, arjunetin and arjungenin on CYP3A4, CYP2D6 and CYP2C9 enzyme activity in human liver microsomes. *Toxicol Rep.* 2015;2:806–16.
 51. Pawar RS, Bhutani KK. Effect of oleanane triterpenoids from Terminalia arjuna--a cardioprotective drug on the process of respiratory oxyburst. *Phytomedicine Int J Phytother Phytopharm.* 2005;12(5):391–3.
 52. Singh G, Singh AT, Abraham A, Bhat B, Mukherjee A, Verma R, et al. Protective effects of Terminalia arjuna against Doxorubicin-induced cardiotoxicity. *J Ethnopharmacol.* 2008;117(1):123–9.
 53. Viswanatha GL shastry, Vaidya SK, C R, Krishnadas N, Rangappa S. Antioxidant and antimutagenic activities of bark extract of Terminalia arjuna. *Asian Pac J Trop Med.* 2010;3(12):965–70.
 54. Ahmad MS, Ahmad S, Gautam B, Arshad M, Afzal M. Terminalia arjuna, a herbal remedy against environmental carcinogenicity: An in vitro and in vivo study. *Egypt J Med Hum Genet.* 2014; 15(1): 61–7.
 55. Mandal S, Patra A, Samanta A, Roy S, Mandal A, Mahapatra TD, et al. Analysis of phytochemical profile of Terminalia

- arjuna bark extract with antioxidative and antimicrobial properties. *Asian Pac J Trop Biomed.* 2013;3(12):960–6.
56. Khan AS. Trees with Antimicrobial Activities. In: *Medicinally Important Trees* [Internet]. Cham: Springer International Publishing; 2017 [cited 2021 Nov 13]. p. 85–108. Available:http://link.springer.com/10.1007/978-3-319-56777-8_4
 57. Aneja KR, Sharma C, Joshi R. Antimicrobial activity of Terminalia arjuna Wight & Arn. an ethnomedicinal plant against pathogens causing ear infection. *Braz J Otorhinolaryngol.* 2012;78(1):68–74.
 58. Verma N, Vinayak M. Effect of Terminalia arjuna on antioxidant defense system in cancer. *Mol Biol Rep.* 2009;36(1):159–64.
 59. Gauthaman K, Banerjee SK, Dinda AK, Ghosh CC, Maulik SK. Terminalia arjuna (Roxb.) protects rabbit heart against ischemic-reperfusion injury: role of antioxidant enzymes and heat shock protein. *J Ethnopharmacol.* 2005;96(3):403–9.
 60. Oberoi L, Akiyama T, Lee K-H, Liu SJ. The aqueous extract, not organic extracts, of Terminalia arjuna bark exerts cardioprotective effect on adult ventricular myocytes. *Phytomedicine.* 2011;18(4):259–65.
 61. Kumar GP, Navya K, Ramya EM, Venkataramana M, Anand T, Anilakumar KR. DNA damage protecting and free radical scavenging properties of Terminalia arjuna bark in PC-12 cells and plasmid DNA. *Free radicals and antioxidants.* 2013;3(1):35-9.
 62. Hebbani AV, Vaddi DR, Dd PP, NCh V. Protective effect of Terminalia arjuna against alcohol induced oxidative damage of rat erythrocyte membranes. *J Ayurveda Integr Med.* 2021;12(2):330–9.
 63. Devi RS, Narayan S, Vani G, Shyamala Devi CS. Gastroprotective effect of Terminalia arjuna bark on diclofenac sodium induced gastric ulcer. *Chem Biol Interact.* 2007;167(1):71–83.
 64. Prakash V, Sehgal V kumar, Bajaj VK, Singh H. To Compare the Effects of Terminalia Arjuna with Rosuvastatin on Total Cholesterol and Low Density Lipoprotein Cholesterol. *Int J Med Dent Sci.* 2016;5(1):1056.
 65. Allawadhi P, Khurana A, Sayed N, Kumari P, Godugu C. Isoproterenol-induced cardiac ischemia and fibrosis: Plant-based approaches for intervention. *Phytother Res PTR.* 2018;32(10):1908–32.
 66. Bhatia J, Bhattacharya SK, Mahajan P, Dwivedi S. Effect of Terminalia arjuna on coronary flow—an experimental study. *Indian J Pharmacol.* 1998;30:118.
 67. Verma P, Muneesh RS, Bhutani G. Experimental Evaluation of Terminalia arjuna (Aqueous Extract) on cardiovascular system in comparison to digoxin. *J Dent Med Sci.* 2013;7:48-51.
 68. Sumitra M, Manikandan P, Kumar DA, Arutselvan N, Balakrishna K, Manohar BM, et al. Experimental myocardial necrosis in rats: role of arjunolic acid on platelet aggregation, coagulation and antioxidant status. *Mol Cell Biochem.* 2001;224(1–2):135–42.
 69. Mythili P, Parameswari CS, Dayana J. Phytochemical analysis of the bark extract of Terminalia arjuna and its cardioprotective effect. *Indian J Innov Dev.* 2012;1:40-2.
 70. Tiwari AK, Gode JD, Dubey GP. Effect of Terminalia arjuna on lipid profiles of rabbits fed hypercholesterolemic diet. *International Journal of Crude Drug Research.* 1990;28(1):43-7.
 71. Pathak SR, Upadhyaya L, Singh RN. Effect of Terminalia arjuna on lipid profile of rabbit fed hypercholesterolemic diet. *International Journal of Crude Drug Research.* 1990;28:48-51.
 72. Khanna AK, Chander R, Kapoor NK. Terminalia arjuna: an ayurvedic cardioprotective, regulates lipid metabolism in hyperlipaemic rats. *Phytotherapy Research.* 1996;10(8):663-5.
 73. Ram A, Lauria P, Gupta R, Kumar P, Sharma VN. Hypocholesterolaemic effects of Terminalia arjuna tree bark. *J Ethnopharmacol.* 1997;55(3):165–9.
 74. Subramaniam S, Subramaniam R, Rajapandian S, Uthrapathi S, Gnanamanickam VR, Dubey GP. Anti-Atherogenic Activity of Ethanolic Fraction of Terminalia arjuna Bark on Hypercholesterolemic Rabbits. *Evid-Based Complement Altern Med ECAM.* 2011;487916.
 75. Subramaniam S, Ramachandran S, Uthrapathi S, Gnanamanickam VR, Dubey GP. Anti-hyperlipidemic and antioxidant potential of different fractions of Terminalia arjuna Roxb. bark against PX- 407 induced hyperlipidemia. *Indian J Exp Biol.* 2011;49(4):282–8.

76. Patil RH, Prakash K, Maheshwari VL. Hypolipidemic effect of *Terminalia arjuna* (L.) in experimentally induced hypercholesteremic rats. *Acta Biologica Szegediensis*. 2011;55(2):289-93.
77. Dwivedi S, CHANSOURIA JN, Somani PN, Udupa KN. Effect of *Terminalia arjuna* on ischaemic heart disease. *Alternative medicine*. 1989;3(2):115-22.
78. Verma SK, Bordia A. Effect of *Terminalia arjuna* bark (*arjun chhal*) in patients of congestive heart failure and hypertension. *J Res Educ Indian Med*. 1988;7(31):e6.
79. Bharani A, Ganguly A, Bhargava KD. Salutary effect of *Terminalia Arjuna* in patients with severe refractory heart failure. *Int J Cardiol*. 1995;49(3):191-9.
80. Parmar HS, Panda S, Jatwa R, Kar A. Cardio-protective role of *Terminalia arjuna* bark extract is possibly mediated through alterations in thyroid hormones. *Pharm*. 2006;61(9):793-5.
81. Bittner, Vera A, Michael Szarek, Philip E. Aylward, Deepak L. Bhatt, Rafael Diaz, Jay M. Edelberg, Zlatko Fras, et al. "Effect of Alirocumab on Lipoprotein(a) and Cardiovascular Risk After Acute Coronary Syndrome. *Journal of the American College of Cardiology*. 2020;75(2):133-44. Available: <https://doi.org/10.1016/j.jacc.2019.10.057>.
82. Garg, Mayank, and Sandip Mohale. "Prevalence of Metabolic Obesity Normal Weight (MONW) in Cardiovascular Disease Patients - A Hospital-Based Case Control Study." *Journal of Evolution of Medical and Dental Sciences-Jemds*. 2020;9(34):2427-31. Available: <https://doi.org/10.14260/jemds/2020/528>.
83. Kamble, T. K., Ankita Kapse, Sunil Kumar, Sourya Acharya, and Aiswarya Ghule. "Study of Myocardial Performance Index in Prediabetes and Its Correlation with Other Cardiovascular Risk Factors. *Journal of Evolution of Medical and Dental Sciences-Jemds*. 2020;9(10): 721-25. Available: <https://doi.org/10.14260/jemds/2020/157>.
84. Acharya, Sourya, Samarth Shukla, and Anil Wanjari. "Subclinical Risk Markers for Cardiovascular Disease (CVD) in Metabolically Healthy Obese (MHO) Subjects." *Journal of Clinical and Diagnostic Research*. 2019;13(6): OC1-6. Available: <https://doi.org/10.7860/JCDR/2019/41317.12890>.
85. Goodman, Shaun G, Philip E. Aylward, Michael Szarek, Vakhtang Chumburidze, Deepak L. Bhatt, Vera A. Bittner, Rafael Diaz, et al. "Effects of Alirocumab on Cardiovascular Events After Coronary Bypass Surgery." *Journal of the American College of Cardiology*. 2019;74(9):1177-86. Available: <https://doi.org/10.1016/j.jacc.2019.07.015>.
86. Singh A, Lodha P, Sharma A. *Terminalia arjuna* Leaf Gall: The Possible Treatment for Sickle Cell Anaemia", *Journal of Pharmaceutical Research International*. 2021;32(41):64-75. DOI: 10.9734/jpri/2020/v32i4131044.

© 2021 Verma and Jogdand; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/80456>