



Antidiarrhoea Activities of Aqueous Extract of *Myrianthus arboreus* Leaves in Albino Rats

N. Nwachoko^{1*}, E. B. Essien² and E. O. Ayalogu²

¹*Department of Biochemistry, Rivers State University, Nkpolu-Oroworukwo, Port Harcourt P.M.B. 5080, Rivers State, Nigeria.*

²*Department of Biochemistry, University of Port Harcourt, Choba, Rivers State, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. Author NN, designed the study, carried out the bench work, wrote the first draft of the manuscript and performed the statistical analysis. Author EBE designed the study, wrote the protocol, supervised, read and approved the final manuscript. Author EOA designed the study, supervised, managed the literature searches, read and approved the final manuscript and supervised the work. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJBCRR/2019/v28i130135

Editor(s):

(1) Dr. K. V. Ramanath, Professor, Department of Pharmacy Practice, SAC College of Pharmacy, B. G. Nagar, Karnataka, India.

Reviewers:

(1) Ene Paschal Chuka, Rivers State University of Science and Technology, Nigeria.
(2) Md. Shariful Islam, Southeast University, Bangladesh.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/46736>

Received 15 November 2018

Accepted 19 January 2019

Published 31 October 2019

Original Research Article

ABSTRACT

Diarrhoea is one of the leading causes of morbidity and mortality in children under the age of 5 years. Due to this problem, the World Health Organization has encouraged studies that will bring about the desired treatment and prevention of diarrhoea. *Myrianthus arboreus* leaves (MA) is used in some tribes of Nigeria for food. In this study, the antidiarrhoea activities of the aqueous extract of *Myrianthus arboreus* leaves were investigated with experimental animals via faecal count, measurement gastrointestinal charcoal meal distance and electrolyte composition. The extract (500, 1000 and 2000) mg/kg in comparison with loperamide hydrochloride, decreased the degree of gastrointestinal motility, production of diarrhoea stool, reduced the frequency of defecation and delayed the onset of diarrhoea in castor oil induced in albino rats. Also the extract inhibited the concentration of intestinal fluid electrolytes.

*Corresponding author: E-mail: blessedconfidence@yahoo.com;

Keywords: *Diarrhoea; Myrianthus arboreu; castor oil; loperamide hydrochloride.*

1. INTRODUCTION

Myrianthus arboreus is a tropical tree growing up to 15m height which contain both male and female flower with spreading branches from a short stem [1-3]. It is usually divided close to the base, the roots form a network structure above ground. It is found in wet environment and on stream-banks of the forest region of Guinea to West Camerouns and extending across Africa. The wood is yellowish-white, soft, fibrous and difficult for art work [4,5]. Though perishable, it is used in making soap. The slash is slightly tinted but rapidly darkens to brown. The bark is said to be variable in appearance: In parts of South East Nigeria, it may be greenish white and slightly flaky, or almost white and smooth [6,5]. In Congo, a bark-tisane is said to be cholagogic and antidyserteric. The bark is taken in Nigeria as a febrifuge. In Nigeria, it is known by Igbo as *Ujuju*, Edo *Ihioghe*, Efik *Ndisok*, Igala *Apulu*, Nupe *Tsakupaci* and Yoruba *Ibi cere* [2,3]. When the leaves fall and lie on the ground, they form a good ground-cover retaining moisture and rotting down to form thick humus. In SE Nigeria, the young leaves are commonly eaten in vegetable-soup, hence the tree being known in Yoruba as the "Soup tree". The soup is so highly considered by the Egba people as to evoke the saying that "one will kill his child for the sake of *ibishere* soup" [2,3]. In Ghana, there are some tribes that eat the leaves as part of their food. In Gabon scrapings of the leaves are cooked in palm-oil is taken to relieve sore-throat. Extract of the leaves is made in Nigeria with *Alehornea* (Euphorbiaceae) for drinking in case of dysentery, and leafy shoots are chewed by people on the Cameroon Mountain for this purpose [6,3]. In Sierra Leone, the decoction form boiled young leaf-flushes and peeled green banana have been used as medicine taken to stop diarrhoea and vomiting. Nwachoko [7] reported that *Myrianthus arboreus* is rich in phytochemicals (glycosides, saponins, sterols and tannins). Amata [8], and Oyeyemi [9] reported that *Myrianthus arboreus* leaves is rich in phytochemicals, noting the presence of tannins, alkaloids and flavonoids. Diarrhoea is the passage of abnormal liquid or unformed stool at an increased frequency [10-12]. It is also linked with increased frequency, fluidity or volume of bowel movements and is characterized by increased frequency of bowel sound movement, wet stool and abdominal pain [13,14]. The passage of loose or watery stools is

usually at least three times in a 24 hour period [14]. In Nigeria, diarrhoea resulting from infection is one of the known killer diseases among children under 5 years of age [15,16]. This work examine the antidiarrhoea activities of aqueous extract of *Myrianthus arboreus* leaves in albino rats.

2. MATERIALS AND METHODS

2.1 Plant Material

Myrianthus arboreus leaves (MA) were obtained from Ekuku-Agbor in Ika South Local Government of Delta State, Nigeria. The sample was air dried and ground to powder form prior to analysis. Extraction was carried out with warm distilled water in the ratio of 1 g to 5 ml. The powdered form of the leaves were soaked for 30 minutes and filtered using Whatman No. 1 filter paper and also with a funnel plunged with glass wool [17]. The residue was re-extracted in warm distilled water for same duration and filtered. The filtrates were pooled together and concentrated with a water bath at 50°C. The concentrated form of the sample was stored at 4°C until needed.

2.2 Experimental Animals

Albino rats weighing between 150 – 200 g, were obtained from the Animal House of the Department of Biochemistry, University of Port Harcourt, Choba, Nigeria. The animals were acclimatized for one week prior to the commencement of the experiment. The animals were housed under standard laboratory conditions with light and dark cycles of 12 hours and were provided with rodent pellet food and water *ad libitum*.

2.3 Drugs

The drugs used in this study included castor oil (finest cold drawn castor oil), loperamide hydrochloride (Aaron Healthcare and Export PVT Ltd, Uttarahand, India), activated choarcoal and gum acacia (Sigma, USA).

2.4 Castor Oil Induced Gastrointestinal Transit

Rats of either sex of weight (150 – 200 g), fasted for 18 hours were randomly allocated into six groups. Group 1 (control) received 10 ml/kg of

Table 1. Effect of aqueous extract of MA leaves on castor oil-induced diarrhoea and Na⁺/K⁺ concentrations in albino rats

Group	Treatment	IL (cm)	CML(cm)	PI(%)	I(%)	Na ⁺ (mmol/l)	K ⁺ (mmol/l)
1	Control	84.0 ± 7.0	47.3 ± 2.5	56.3	-	25.7±4.6 ^a	6.3±3.2 ^a
2	Untreated	86.3 ± 3.8	21.0 ± 7.5	24.3	-	10.0±0.0a*	3.6±0.5 ^{a*}
3	500mg/kg	77.7 ± 6.1	30.3 ± 9.3	39.0	35.9	10.0±0.0 ^a	2.9±0.4 ^a
4	1000mg/kg	83.3 ± 4.2	32.7 ± 4.6	39.0	30.9	12.0±2.0 ^a	3.7±0.4 ^a
5	2000mg/kg	95.3 ± 5.0	32.7 ± 4.7	34.3	30.9	16.3±6.0 ^a	2.8±0.2 ^a
6	5mg/kg loperamide	89.0 ± 8.9	27.7 ± 9.7	31.1	41.4	18.7±15.0 ^a	4.6±1.7 ^a

Key; LSI – length of small intestine, CML = Charcoal meal length, PI = Peristaltic index,
I = Inhibition, MA = Myrianthus arboreus. n= 4

Table 2. Effect of aqueous extract of MA on the faecal count of castor oil-induced diarrhoea albino rats

Group	Treatment	OD (MIN)	MWF	% I
1	Control	30	1.5 ± 2.0	-
2	500 mg/kg	240	0.3 ± 0.0	80.0
4	1000 mg/kg	60	0.7 ± 1.4	53.3
4	2000 mg/kg	240	0.2 ± 0.0	89.0
5	5 mg/kg loperamide	240	0.3 ± 0.0	80.0

Key; OD = Onset of diarrhoea, MWF = Mean wet faeces after 6 hour,
I = Inhibition

distilled water, group 2 (normal) received no form of pre-treatment, group 3, 4 and 5 received 500, 1000 and 2000 mg/kg of MA extract and group 6 received 5mg/kg of loperamide hydrochloride. After 1 hour of treatment with the extract, distilled water and standard drug, diarrhoea was induced by oral administration of 1ml of castor oil to the test as well as the control group, after a period of 1 hour latter, 1 ml of charcoal meal (10% charcoal suspension and 5% gum acacia) was given to each of the animals. The animals were sacrificed after 1 hour following the administration charcoal meal. The distance traveled by the charcoal meal from the pylorus to the caecum were measured and expressed as a percentage of the total length of the intestine from the pylorus to the caecum of each animal [18-20].

PI = LM/LSI x 100%.

PI = Peristaltic index,

LM – Distance travelled by charcoal meal,

LSI – length of small intestine, % inhibition: (control – test)/control x 100.

2.5 Castor Oil Induced Diarrhoea in Rats and Faecal Count

Castor oil-induced diarrhoea was determined by the method of [21] Awoutes et al. (1978). Rats weighing between 150 – 200 g fasted for 18 hours were randomly distributed into five groups. The administration procedure was as described above. The time taken for onset of diarrhoea and faecal droppings were recorded. Percentage inhibition was calculated [22,19,23].

% inhibition = (control – test)/control x 100.

3. RESULTS

The effect of aqueous extract of MA leaves on castor oil-induced diarrhoea and Na⁺/K⁺ concentrations in albino rats is as shown in Table 1.

The effect of aqueous extract of MA on the faecal count of castor oil-induced diarrhoea albino rats is as shown in Table 2.

4. DISCUSSION

The result of the percentage inhibition of the aqueous extract of *Myrianthus arboreus* leaves (MA) against castor oil induced diarrhoea in albino rats are as shown in Table 1. The result showed that mean charcoal meal length (CML) of

animals in group 1 was the highest, followed by the mean CML of animals in group 4 and 5 pretreated with 1000 and 2000 mg/kg MA extract respectively. Trend observed in the mean CML of animals in a group will reflect in the percentage peristaltic index (PI) and percentage inhibition. Thus group 6 animals with the least mean CML had the least PI and the highest percentage inhibition. Also group 4 and 5 animals that have the highest CML among the pretreated groups, have the highest PI and least percentage inhibition. Comparing the PI values of both control and the treated groups, MA leaf extract inhibited the PI values of treated groups. Also, the distance travelled by charcoal meal in the experimental groups was inhibited by the extract. The distance traveled by charcoal meal in the control, untreated and treated groups, showed that MA leaf possess antidiarrhoea properties. Also the extract may have affected the concentration of intestinal fluid electrolytes. Na⁺ value for group 1 (control) was 25.7 ± 4.6 mmol/l, which was higher when compared with that of the treated groups.

Table 2 showed the result of percentage inhibition of aqueous extract of MA leaves on the faecal count of castor oil induced diarrhoea in albino rats and the delay to the onset of diarrhoea. The results show that the control group (group1) had minimal resistance to the onset of diarrhoea. Following 30 minutes after the administration of castor oil, the animals in the group produced wet/diarrhoea stool. Different is the case of the groups pretreated with MA extract and standard drug. The animals in group 3 had 60 minutes delay while groups 2, 4 and 5 had 240 minutes delay respectively, before the onset of diarrhoea. The same trend was observed in the mean wet faeces (MWF) in both control and the pretreated groups. Group 1 animals which received no form of pretreatment had 1.5 ± 2.0 MWF while the animals in group 3 with a delay of 60 minutes, had MWF of 0.7 ± 1.4. The animals in groups 2 and 5 had MWF of 0.3 ± 0.0 each. Groups that had the highest delay in the onset of diarrhoea, had the least MWF and highest percentage inhibition.

4. CONCLUSION

In conclusion, it could be said that, on evaluating the antidiarrhoea property of MA leaves via inhibition of distance traveled by charcoal meal, wet faecal count and intestinal fluid electrolytes, it could be said that MA leaves extract possess significant antidiarrhoea property.

ETHICAL APPROVAL

As per international standard or university standard ethical approval has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- White L, Abernathy K. A guide to the vegetation of the Lopé Reserve, Gabon. Wildlife Conservation Society, New York, United States. 1997;224.
- Wilks C, Issembe Y. Les arbres de la Guinée Equatoriale: Guide pratique d'identification: région continentale. Projet CUREF, Bata, Guinée Equatoriale. 2000; 546 .
- Tshibangu JN, Chifundera K, Kaminsky R, Wright AD, Konig GM. Screening of African medicinal plants for antimicrobial and enzyme inhibitory activity. Journal of Ethnopharmacology. 2002;80:25–35.
- Irvine FR. Woody plants of Ghana, with special reference to their uses; 1961.
- Okigbo BN. Vegetables in tropical Africa. In: Crop genetic resources in Africa. Proceedings of a workshop jointly organized by the Association for the Advancement of Agricultural Sciences in Africa and the International. Institute of Tropical Agriculture held at IITA Ibadan, Nigeria. 1978;128–147.
- Okafor JC (). Development of forest tree crops for food supplies in Nigeria. Forest Ecology and Management. 1978;1:235–247.
- Nwachoko N, Essien EB, Ayalogu EO. Proximate and quantitative phytochemical analysis of *Myrianthus arboreus*. Global Institute for Research, Conference Book of Proceedings. 2015;3(1):81-85.
- Amata AI. Nutritive value of the leaves of *Myrianthus arboreus*. A Brownse plant. International Journal of Agricultural Research. 2010;5:576-581.
- Oyeyemi SD, Arowosegbe S, Adebisi AO. Phytochemical and proximate evaluation of *Myrianthus arboreus*. Journal of Agriculture and Veterinary Science. 2014;7(9):1-5.
- Guerrant RL, Schorling JB, McAuliffe JF, De Souza MA. Diarrhea as a cause and an effect of malnutrition: diarrhea prevents catch-up growth and malnutrition increases diarrhea frequency and duration. The American Journal of Tropical Medicine and Hygiene. 1992;47(1 Pt 2):28–35.
- Ahlquist DA. Constipation and diarrhoea. In: Principles of internal medicine. McGraw Hill Medical Publishing Division, New York, 2001;241-247.
- Wilson ME. Diarrhea in non travelers: Risk and etiology. Clinical Infectious Disease. 2005;41(8):51–56.
- WHO. Diarrhoea. Wikipedia the free encyclopedia; 2003.
- Navaneethan U, Giannella RA. Mechanisms of infectious diarrhea. Nature Clinical Practice Gastroenterology and Hepatology. 2008;5(11):637–47.
- Audu R, Umilabug SA, Renner JK, Awodiji JA. Diarrhoea management. Journal of Nigerian Infection Control Association. 2000;3:15.
- Nwachoko N, Jack IR. Phytochemical screening and antidiarrhoea activities of *Tetracarpidium conophorum* induce in albino rats. Sky Journal of Biochemistry Research. 2015;4(3):16–20.
- Bakare RI, Magbagbeola OA, Akinwande AI, Okunowo OW, Green M. Anti-diarrhoeal activity of aqueous leaf extract of *Momordica charantia* in rats. Journal of Pharmacognosy and Phototherapy. 2011; 3(1):1-7.
- Mascolo N, Izzo AA, Antore FB, Capasso F. Nitric oxide and castor oil induced diarrhea. Journal of Pharmacological Experiment. 1994;128:291– 293.
- Mukherjee PK, Saha K, Murugeban T. Screening of Antidiarrhoea Profile of Some Plant Extracts of a Specific Region of West Bengal, India. Journal of Ethnopharmacology. 1998;60:85–89.
- Rani SN, Ahamed N, Rajaram S, Saluja R. Atidiarrhoea evaluation of *Clerodendrum phlomides* linn leaf extract in rats. Journal of Ethno-Pharmacology. 1999;68:315–319.
- Awouters F, Niemegeers CJE, Lenaerts FM, Janseen PA. Delay of castor oil diarrhoea in rats: A new way to evaluate inhibitors of

- prostaglandin biosynthesis. Journal of Pharmaceutical Pharmacology. 1978;30: 4–45.
22. Izzo AA, Nicholetti M, Giamathasis B, Capasso F. Antidiarrhoea activity of *Fermentia servicea* Burchex DC Extract in: Capasse, F, Mascolo, N. (Eds), National Drug and Digestive Tract. EMSI, Rome, 1992;223–230
23. Karim AH, Maklfia A, Zigyat AL, Bnouham M. Antidiarrhoea activity of crude aqueous extract of *Rubia Tinctorum* L. Roots in rident. Journal Smooth Muscle Resource. 2010;46(2):119–123.

© 2019 Nwachoko et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/46736>